

# Improving cognition and healthy lifestyle behavior in chronic obstructive pulmonary disease

## Citation for published version (APA):

van Beers, M. C. M. (2021). *Improving cognition and healthy lifestyle behavior in chronic obstructive pulmonary disease*. Maastricht University. <https://doi.org/10.26481/dis.20210421mb>

## Document status and date:

Published: 01/01/2021

## DOI:

[10.26481/dis.20210421mb](https://doi.org/10.26481/dis.20210421mb)

## Document Version:

Publisher's PDF, also known as Version of record

## Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

## General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.umlib.nl/taverne-license](http://www.umlib.nl/taverne-license)

## Take down policy

If you believe that this document breaches copyright please contact us at:

[repository@maastrichtuniversity.nl](mailto:repository@maastrichtuniversity.nl)

providing details and we will investigate your claim.

# Improving cognition and healthy lifestyle behavior in chronic obstructive pulmonary disease

## Citation for published version (APA):

van Beers, M. C. M. (2021). *Improving cognition and healthy lifestyle behavior in chronic obstructive pulmonary disease*. Maastricht University. <https://doi.org/10.26481/dis.20210421mb>

## Document status and date:

Published: 01/01/2021

## DOI:

[10.26481/dis.20210421mb](https://doi.org/10.26481/dis.20210421mb)

## Document Version:

Publisher's PDF, also known as Version of record

## Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

## General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.umlib.nl/taverne-license](http://www.umlib.nl/taverne-license)

## Take down policy

If you believe that this document breaches copyright please contact us at:

[repository@maastrichtuniversity.nl](mailto:repository@maastrichtuniversity.nl)

providing details and we will investigate your claim.

Download date: 26 Jul. 2021

# **Improving cognition and healthy lifestyle behavior in chronic obstructive pulmonary disease**

Martinus Cornelius Maria van Beers



© Copyright Martijn van Beers, Maastricht, 2021

Cover design: Frits Roothans

Layout: Martijn van Beers

Printed by: ProefschriftMaken || [www.proefschriftmaken.nl](http://www.proefschriftmaken.nl)

ISBN: 978-94-6423-103-8

The research described in this thesis was performed at the Department of Respiratory Medicine, NUTRIM School of Nutrition and Translational Research in Metabolism at Maastricht University Medical Center+, Maastricht, The Netherlands

The research was financially supported by grants from ZonMW (JPI-HDHL 'Ambrosiac' grant, project number 529051006) and the Maastricht University Interfaculty Program 'Eatwell'

# **Improving cognition and healthy lifestyle behavior in chronic obstructive pulmonary disease**

Proefschrift

Ter verkrijging van de graad van doctor aan de Universiteit Maastricht,  
op het gezag van de Rector Magnificus, prof. dr. Rianne M. Letschert,  
volgens het besluit van het College van Decanen,  
in het openbaar te verdedigen op donderdag  
11 februari 2021 om 13:00 uur

door

Martinus Cornelius Maria van Beers

Geboren op 3 juni 1994 te Veldhoven, Nederland

**Promotor**

Prof. dr. A. M. W. J. Schols

**Co-promotor**

Dr. H. R. Gosker

**Beoordelingscommissie**

Prof. dr. A. Jansen (voorzitter)

Prof. dr. L. C. P. G. M. de Groot (Wageningen University)

Dr. S. Köhler

Prof. dr. D. van de Mheen (Tilburg University)

Prof. dr. G. Wesseling

# Table of Contents

Chapter 1	General Introduction	7
Chapter 2	Clinical outcome and cost-effectiveness of a 1-year nutritional intervention program in COPD patients with low muscle mass: the randomized controlled NUTRAIN trial	25
Chapter 3	Motivational regulations and lifestyle in patients with COPD and low muscle mass before and after pulmonary rehabilitation	61
Chapter 4	Cognitive impairment in chronic obstructive pulmonary disease: disease burden, determinants and possible future interventions	85
Chapter 5	Cognitive performance in relation to metabolic disturbances in patients with COPD	121
Chapter 6	Feasibility and efficacy of working memory training in chronic obstructive pulmonary disease: the randomized placebo-controlled Cogtrain trial	145
Chapter 7	General Discussion	187
	Summary	207
	Samenvatting	213
	Impact	219
	Dankwoord	227
	Curriculum vitae	233
	List of publications	237





# Chapter 1

General Introduction

## Chronic obstructive pulmonary disease

Chronic obstructive pulmonary disease (COPD) is a common and preventable chronic disease which is characterized by airflow limitation that is usually progressive (1). An estimated 299 million people worldwide suffered from the disease in 2017 (2) and it caused 3 million deaths in 2016, making it the third most common cause of death worldwide (3). Tobacco smoking and exposure to air pollution, toxic gases and biomass fuel emission are the most common risk factors for developing COPD (4).

The pathophysiology in the lungs primarily consists of chronic bronchitis (airway inflammation) and emphysema (destruction of the small airways) (1), but COPD is highly heterogeneous. Various systemic manifestations outside the lungs and airways are common. These include cardiovascular disease (5), musculoskeletal impairment (6), cachexia (7), sarcopenia (8), metabolic syndrome (MetS) (9), anxiety and depression (10, 11) and cognitive abnormalities (e.g., 12, 13). The entire constellation of COPD manifestations contributes to increased disease burden, worsened health status and functional performance, increased hospitalization and mortality risk and higher healthcare costs (14-16).

COPD is incurable but can be treated. Smoking cessation is the most effective prevention as well as treatment strategy. The current guidelines recommend additional pharmacological treatment using a combination of long-acting muscarinic antagonists and  $\beta_2$ -agonists, combined in severe cases with inhaled corticosteroids (1). Patients who remain symptomatic despite optimal treatment and/or those with complex pathology (such as poor lung function, high burden of comorbidities, a poor functional or nutritional status and/or frequent exacerbations) can be referred to pulmonary rehabilitation (PR) (17) or in very specific cases be eligible for lung volume-reducing endoscopic or surgical intervention (1).

## Pulmonary rehabilitation

PR is defined as "a comprehensive intervention based on a thorough patient assessment followed by patient-tailored therapies that include, but are not limited to, exercise training, education and behavior change, designed to improve the physical and psychological condition of people with chronic respiratory disease and to promote the long-term adherence to health-enhancing behaviors" (17).

PR is one of the most effective interventions to improve the health status of patients with COPD. Exercise capacity and health-related quality of life can

increase significantly over the course of a PR program (18). However, the long-term efficacy of PR has not been investigated yet. This is relevant because PR-induced gains translate only poorly into a healthier lifestyle, such as more physical activity, after PR (18). Moreover, specific subgroups of patients could benefit from specific additional treatments, such as nutritional supplementation (19, 20). These considerations were the main rationale to design the randomized, double-blind, placebo-controlled NUTRAIN trial. This trial investigated the efficacy and cost-effectiveness of a multimodal nutritional management strategy, consisting of nutritional supplementation and dietary counseling, in COPD patients with low muscle mass during a 4-month outpatient PR program and in the 8-month maintenance phase thereafter. The results of the initial 4-month phase were published earlier (21); the efficacy and cost-effectiveness of the overall strategy are reported in Chapter 2 of this thesis.

Nutritional supplementation did not improve quadriceps muscle strength during the initial PR phase of the NUTRAIN trial. Interestingly, however, supplementation prevented declines in daily physical activity (21). This dissociation between physical capacity and activity, combined with positive effects on plasma micronutrient status and quality of life (21) led to the hypothesis that mechanisms affecting cognitive performance and mental health might be involved.

The main aim of this thesis was therefore to investigate determinants of cognitive performance in patients with COPD and the potential efficacy of working memory training (WMT) next to motivational regulations in increasing patients' adherence to a healthy lifestyle.

## **Motivational regulations as determinants of PR-induced health improvement**

The central tenet of self-determination theory (SDT) is that motivation is the key determinant of behavior (22). SDT conceptualizes motivation as a quantitative construct (i.e., one can be more or less motivated towards something), but it also distinguishes six qualitatively different motivational regulations which can differentially drive behavior. These regulations can be placed on a continuum, ranging from being more intrinsic (performing a behavior because of the enjoyment derived from doing it) to more extrinsic forms of motivation (performing a behavior because of some secondary rewards associated with it) (23). Although all motivational regulations can drive behavior, a larger reliance on more intrinsic regulations as opposed to more extrinsic ones is generally considered beneficial.

1 Intrinsic behavioral regulation is more stable and enduring and benefits one's wellbeing more compared to controlled regulation (24).

Because of their important role in determining behavior, motivational regulations could be the key to ensure PR-induced gains translate into a longer-term healthy lifestyle after the PR program is completed and the associated external support is not present anymore. If motivational regulations are related to healthy lifestyle behaviors before, during and/or after a PR program, they could be used as predictors of adherence to healthy lifestyle behaviors, or even as interventional targets to improve this adherence. Earlier studies have already shown that motivational interviewing, which is a psychological counseling technique based on SDT, can increase medication adherence (25, 26), exercise compliance (26) and health-related quality of life (26-28) and can reduce hospitalization rates (26, 27) in patients with COPD.

Chapter 3 investigates the value of motivational regulations to predict the carry-over of PR-induced healthy lifestyle gains into a healthier lifestyle.

## Cognitive functioning

Cognitive functioning involves any brain function required for perceiving, registering, storing, retrieving and using information from the environment in order to be able to adapt one's behavior to novel situations (29). Cognitive functioning is not a unitary construct but consists of different functions, such as memory, attention, self-control, task switching, inhibition and many more. Furthermore, 'lower' functions can be distinguished from 'higher' ones, with the lower functions (e.g., sensory processing, attention and language abilities) serving as the foundation for the higher ones (e.g., cognitive flexibility, self-control and problem-solving) (29).

Executive functions encompass various higher-order cognitive functions involved in behavioral control. Specifically, inhibition, task switching and working memory (WM) are generally considered to jointly constitute executive functioning (30). Inhibition refers to the ability to override or interrupt undesired behavioral responses (31), and enables to focus on one particular thing instead of being distracted by other items in our environment. Task switching is the ability to quickly switch one's focus from one item to another, and to change demands and priorities (30). This ability makes it possible to avoid behavioral interference from previous tasks, thereby preventing perseveration when a behavioral strategy does not lead to attaining a certain goal (anymore) (32). Finally, WM refers to the ability

to temporarily keep important information in mind, enabling the information to be mentally manipulated (33). WM is also involved in the ability to focus one's attention (34) and to actively keep a representation of long-term goals (35).

There are large inter-individual differences in cognitive functioning. One person's WM might be very strong (for instance if he or she is able to remember and immediately reproduce an entire phone number after hearing it once), whereas in someone else it might be very poor. Likewise, there are large intra-individual differences in cognitive functioning. Someone might simultaneously have a very strong long-term memory but very poor inhibition skills.

### *Measuring cognitive performance*

Cognitive performance can be quantified by using standardized cognitive tests. Because so many different cognitive domains exist, countless tests have been developed to gauge people's performance on those domains. Cognitive tests can be distinguished into screening tools and comprehensive neuropsychological assessment batteries (36).

Screening tools, such as the Mini-Mental State Examination (MMSE) (37), the Montreal Cognitive Assessment (MoCA) (38) and the Addenbrooke's Cognitive Examination-Revised (ACE-R) (39), can be administered relatively quickly (often within 10 to 15 minutes), are generally narrow in scope, and they do not have a diagnostic purpose (36). Screening tools are used in clinical practice to obtain a quick overview of the cognitive performance of a person presenting with memory problems and to determine whether he or she can benefit from referral for more extensive cognitive assessment.

More detailed assessment is conducted using comprehensive neuropsychological test batteries (36), such as the Cambridge Neuropsychological Test Automated Battery (CANTAB). It takes more time and effort to administer such batteries, but their broader scope (i.e., their ability to measure more cognitive domains more thoroughly) and their higher sensitivity to detect CI are large advantages compared to screening tools (36). Therefore, such batteries can be used for diagnostic purposes in a clinical setting (36) and as valid and reliable outcome measures for clinical trials in a scientific setting (40).

## **Cognitive impairment in COPD and its consequences**

The large impact of COPD on the brain and cognitive functioning has long been under-recognized, but the awareness for the neuropsychological consequences

1 of COPD has increased in recent years (e.g., 12, 13). The prevalence of CI is significantly higher in patients with COPD compared to non-COPD controls. A recent large review reported a 32% prevalence of any CI and a 25% prevalence of mild CI (MCI) in patients with COPD (41). More recently, a 39.4% prevalence of CI was reported in clinically stable patients (42), and a 41.5% (43) and 56.7% (13) prevalence in patients referred to PR. In contrast, the prevalence of CI among non-COPD controls in the latter study was 13.3%. In another study in a PR context, 9.5% of patients with COPD scored below the 5th percentile of a normal reference group on the MoCA, and only 38.1% of patients scored higher than the median of the reference group (44).

The consequences of CI extend beyond the cognitive domain. CI affects self-management skills, which are especially relevant in patients with chronic diseases. Cognitive functioning is related to the accuracy of recalling severity of fatigue and dyspnea over a two-week period (45) and to inhaler competency (46) in patients with COPD. Impaired prospective memory and planning can lead to decreased treatment and medication adherence (47). The risks of hospitalization and mortality (48) are higher in patients with CI compared to those without. Cognitive functioning is also related to engagement in unhealthy lifestyle behaviors, such as food intake in overweight people (49) and alcohol consumption in problem drinkers (50). The relationship between CI and health-related quality of life remains unclear: in some studies the two were significantly negatively related (42, 51, 52) but not in others (43, 53).

Cognitive stress susceptibility is already heightened by chronic disease (54), and CI and cerebral abnormalities might further amplify these effects: executive functioning is related to cognitive stress reactivity (55) and perception (56), and the prefrontal cortex and hippocampus are smaller in patients with COPD compared to healthy controls (57-59). These are the two main areas involved in the regulation of the hypothalamic-pituitary-adrenal (HPA) axis, which is activated in response to stress (60, 61). Moreover, exposure to cortisol, which is the end-product of HPA axis activation, contributes to shrinkage of these two areas (62-64), thereby creating a vicious cycle of progressively poorer stress regulation.

### *Potential etiological factors*

Many typical or common factors associated with COPD are independently related to poorer cognitive functioning. All of these likely contribute to CI in COPD, depending on the extent to which they affect any given patient. Disease-specific factors include inflammation (65, 66), exacerbations (51, 67, 68), hypoxemia and hypercapnia (69, 70). Obstructive sleep apnea (71, 72), chronic heart failure (41)

and depression (73) are common comorbidities of COPD. In addition, modifiable lifestyle factors exhibited by many patients with COPD, such as a history of smoking (74), physical inactivity (75-77) and a low-quality diet (78) play a role. Chapter 4 of this dissertation reviews the initial knowledge base around the specific cognitive domains which are predominantly affected in COPD, the prevalence and pathophysiology of CI in COPD and the impact of CI on health, self-management and daily life outcomes.

Metabolic abnormalities, such as MetS and sarcopenia, are more common in COPD compared to non-COPD controls (8, 9), and both are independently related to poorer cognitive performance (79-83). It is still unknown, however, whether MetS and sarcopenia amplify the detrimental impact of COPD on cognitive performance. If MetS and sarcopenia are related to CI in COPD, they could constitute prognostic and potential screening factors for the development of CI and/or interventional targets to prevent or counteract its development. Therefore, Chapter 5 investigates in patients with COPD referred for PR whether cognitive performance is poorer in patients with MetS or sarcopenia compared to those without.

## **Treating cognitive impairment in COPD**

The large impact of CI on the disease management, health outcomes, lifestyle behaviors and cognitive stress susceptibility of patients with COPD emphasizes the importance to treat CI. Cognitive performance is trainable: the brain is plastic (84, 85). Cognitive performance can therefore be improved through repeated practice. One earlier study investigated the effects of cognitive training in patients with COPD, which did not find improved cognitive performance (86). However, this intervention had a relatively low training load and was aimed at improving attention and logical-deductive thinking. In contrast, WM training (WMT) would be especially promising in COPD, because WM is the principal cognitive domain underlying both general cognitive performance (87) and top-down behavioral regulation.

The connection between WM and general cognitive performance implies that the likelihood of training-induced cognitive improvement 'spilling over' into other cognitive domains is largest when WM is trained, as opposed to other cognitive domains. This 'spill-over' is known in the neuropsychological literature as 'transfer' (88). Most cognitive or WM training interventions are aimed at children or adolescents, because their brains are more plastic than those of the elderly (89).

1 Studies that specifically investigate WMT in healthy elderly populations yield mixed results (90-92), but WMT has never been administered to patients with COPD yet.

The connection between WM and top-down behavioral control is explained by dual-process theories of cognitive functioning (93-95). These postulate the existence of two cognitive processing systems: a fast system and a slower one (95, 96). The fast system is responsible for executing relatively easy, well-known and/or highly automated cognitive processes; the slow system for more deliberate, controlled and conscious execution of harder cognitive tasks (96). WM underlies the latter system (97), which is responsible for top-down behavioral control. Strengthening it would make it easier to execute controlled responses compared to less opportune automatic ones. Several studies have confirmed that cognitive training can decrease engagement in unhealthy lifestyle behaviors, such as overeating in obese people (49) and consuming alcohol in problem drinkers (50), but the effects of cognitive training or WMT on adherence to healthy lifestyle behaviors such as a healthy diet or physical activity (PA) have remained under-investigated. This is relevant in COPD because a sedentary lifestyle is common in advanced stages of COPD (98-100), as is a poor dietary quality. In 564 patients referred for PR, protein intake deviated from the recommended daily intake in 30% of patients, carbohydrate intake in 26% and dietary fat intake in almost one third (101). Intake of many micronutrients, such as iron, sodium, zinc, niacin, folic acid (102), vitamin B6, B9 and B12,  $\beta$ -carotene (103), vitamins A, C, D (101) and E (101, 103) was low.

Chapter 6 therefore presents the results of the randomized, placebo-controlled Cogtrain trial, investigating whether a 12-week home-based WMT program followed by a 12-week maintenance program could establish and maintain improvement in cognitive performance, physical capacity and activity, dietary quality, cognitive stress susceptibility and perception, recall of prespecified healthy lifestyle goals, healthy lifestyle motivation and psychological wellbeing in patients with COPD.

The results of the current thesis will be discussed and placed in a broader context in Chapter 7, with a particular focus on cognitive functioning in COPD and on the potential efficacy of WMT and other promising interventions to improve cognitive functioning and adherence to a healthy lifestyle in patients with COPD.



## References

1. Global Initiative for Chronic Obstructive Lung Disease (GOLD). Global Strategy for the Diagnosis, Management and Prevention of Chronic Obstructive Pulmonary Disease. Agusti A, Vogelmeier C, editors: 2020.
2. GBD 2017 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392(10159):1789-858.
3. World Health Organization. The top 10 causes of death 2018 [accessed July 16, 2020]. Available from: <https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death>.
4. Rabe KF, Watz H. Chronic obstructive pulmonary disease. *Lancet*. 2017;389(10082):1931-40.
5. Chen W, Thomas J, Sadatsafavi M, Fitzgerald JM. Risk of cardiovascular comorbidity in patients with chronic obstructive pulmonary disease: a systematic review and meta-analysis. *Lancet Respiratory Medicine*. 2015;3(8):631-9.
6. Van den Borst B, Slot IGM, Hellwig VACV, Vosse BAH, Kelders MCJM, Barreiro E, *et al*. Loss of quadriceps muscle oxidative phenotype and decreased endurance in patients with mild-to-moderate COPD. *Journal of Applied Physiology*. 2013;114(9):1319-28.
7. Society on Sarcopenia, Cachexia and Wasting Disorders. Definition of cachexia and sarcopenia [accessed October 3, 2020]. Available from: <https://society-scwd.org/cachexia-definition/>.
8. Trajanoska K, Schoufour JD, Darweesh SKL, Benz E, Medina-Gomez C, Alferink LJM, *et al*. Sarcopenia and its clinical correlates in the general population: the Rotterdam Study. *Journal of Bone and Mineral Research*. 2018;33(7):1209-18.
9. Cebron Lipovec N, Beijers RJ, Van den Borst B, Doehner W, Lainscak M, Schols AMWJ. The prevalence of metabolic syndrome in chronic obstructive pulmonary disease: a systematic review. *COPD*. 2016;13(3):399-406.
10. Ouellette DR, Lavoie KL. Recognition, diagnosis and treatment of cognitive and psychiatric disorders in patients with COPD. *International Journal of COPD*. 2017;12:639-50.
11. Panagioti M, Scott C, Blakemore A, Coventry PA. Overview of the prevalence, impact and management of depression and anxiety in chronic obstructive pulmonary disease. *International Journal of Chronic Obstructive Pulmonary Disease*. 2014;9:1289-306.
12. Dodd JW, Getov SV, Jones PW. Cognitive function in COPD. *European Respiratory Journal*. 2010;35:913-22.
13. Cleutjens FAHM, Franssen F, Spruit MA, Vanfleteren LEGW, Gijzen C, Dijkstra JB, *et al*. Domain-specific cognitive impairment in patients with COPD and control subjects. *International Journal of Chronic Obstructive Pulmonary Disease*. 2017;12:1-11.

14. Sievi N, Senn O, Brack T, Brutsche MH, Frey M, Irani S, *et al.* Impact of comorbidities on physical activity in COPD. *Respirology*. 2015;20(3):413-8.
15. Mannino DM, Thorn D, Swensen A, Holguin F. Prevalence and outcomes of diabetes, hypertension and cardiovascular disease in COPD. *European Respiratory Journal*. 2008;32(4):962-9.
16. Mannino DM, Higuchi K, Yu T-C, Zhou H, Li Y, Tian H, *et al.* Economic burden of COPD in the presence of comorbidities. *Chest*. 2015;148(1):138-50.
17. Spruit MA, Singh SJ, Garvey C, ZuWallack R, Nici L, Rochester C, *et al.* An official American Thoracic Society / European Respiratory Society statement: Key concepts and advances in pulmonary rehabilitation. *American Journal of Respiratory and Critical Care Medicine*. 2013;188(8):e13-e64.
18. Troosters T, Demeyer H, Hornikx M, Camillo CA, Janssens W. Pulmonary rehabilitation. *Clinics in Chest Medicine*. 2014;35(1):241-9.
19. Hornikx M, Van Remoortel H, Lehouck A, Mathieu C, Maes K, Gayan-Ramirez G, *et al.* Vitamin D supplementation during rehabilitation in COPD: a secondary analysis of a randomized trial. *Respiratory Research*. 2012;13:84.
20. Broekhuizen R, Wouters EF, Creutzberg EC, Weling-Scheepers CA, Schols AM. Polyunsaturated fatty acids improve exercise capacity in chronic obstructive pulmonary disease. *Thorax*. 2005;60:376-82.
21. Van de Bool C, Rutten EPA, Van Helvoort A, Franssen FM, Wouters EF, Schols AM. A randomized clinical trial investigating the efficacy of targeted nutrition as adjunct to exercise training in COPD. *Journal of Cachexia, Sarcopenia and Muscle*. 2017;8:748-58.
22. Deci EL, Ryan RM. *Intrinsic motivation and self-determination in human behavior*. New York, NY: Plenum; 1985.
23. Deci EL, Ryan RM. The "what" and "why" of goal pursuits: human needs and the self-determination of behavior. *Psychological Inquiry*. 2000;11:227-68.
24. Markland D, Ryan RM, Tobin VJ, Rollnick S. Motivational interviewing and self-determination theory. *Journal of Social and Clinical Psychology*. 2005;24(6):811-31.
25. Naderloo H, Vafadar Z, Eslaminejad A, Ebadi A. Effects of motivational interviewing on treatment adherence among patients with chronic obstructive pulmonary disease: a randomized controlled clinical trial. *Tanaffos*. 2018;17(4):241-9.
26. Long H, Howells K, Peters S, Blakemore A. Does health coaching improve health-related quality of life and reduce hospital admissions in people with chronic obstructive pulmonary disease? A systematic review and meta-analysis. *British Journal of Health Psychology*. 2019;24:515-46.

27. Benzo RP, Vickers Douglas K, Novotny PJ, Tucker S, Hoult J, Neuenfeldt P, *et al.* Health coaching and chronic obstructive pulmonary disease rehospitalization: a randomized study. *American Journal of Respiratory and Critical Care Medicine*. 2016;194(6):672-80.
28. Rehman H, Karpman C, Douglas KV, Benzo R. Effect of a motivational interviewing-based health coaching on quality of life in subjects with COPD. *Respiratory Care*. 2017;62(8):1043-8.
29. Cleutjens FA, Janssen DJ, Ponds RW, Dijkstra JB, Wouters EF. COgnitive-Pulmonary Disease. *BioMed Research International*. 2014;697825.
30. Diamond A. Executive functions. *Annual Review of Psychology*. 2013;64:135-68.
31. Logan GD, Cowan WB, Davis KA. On the ability to inhibit simple and choice reaction time responses: a model and a method. *Journal of Experimental Psychology: Human Perception and Performance*. 1984;10(2):276-91.
32. Miyake A, Friedman NP, Emerson MJ, Witzki AH, Howerter A, Wager TD. The unity and diversity of executive functions and their contributions to complex "frontal lobe" tasks: a latent variable analysis. *Cognitive Psychology*. 2000;41(1):49-100.
33. Baddeley A. Working memory. *Science*. 1992;255(5044):556-9.
34. Barrett LF, Tugade MM, Engle RW. Individual differences in working memory capacity and dual-process theories of the mind. *Psychological Bulletin*. 2004;130(4):553-73.
35. Hofmann W, Gschwendner T, Friese M, Wiers RW, Schmitt M. Working memory capacity and self-regulatory behavior: Toward an individual differences perspective on behavior determination by automatic versus controlled processes. *Journal of Personality and Social Psychology*. 2008;95(4):962-77.
36. Roebuck-Spencer TM, Glen T, Puente AE, Denney RL, Ruff RM, Hostetter G, *et al.* Cognitive screening tests versus comprehensive neuropsychological test batteries: a National Academy of Neuropsychology education paper. *Archives of Clinical Neuropsychology*. 2017;32(4):491-8.
37. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": a practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*. 1975;12(3):189-98.
38. Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, *et al.* The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *Journal of the American Geriatrics Society*. 2005;53(4):695-9.
39. Mioshi E, Dawson K, Mitchell J, Arnold R, Hodges JR. The Addenbrooke's Cognitive Examination Revised (ACE-R): a brief cognitive test battery for dementia screening. *International Journal of Geriatric Psychiatry*. 2006;21(11):1078-85.
40. Lezak MD, Howieson DB, Bigler ED, Tranel D. *Neuropsychological Assessment* (5th edition). New York, NY: Oxford University Press; 2012.

41. Yohannes AM, Chen W, Moga AM, Leroi I, Connolly MJ. Cognitive impairment in chronic obstructive pulmonary disease and chronic heart failure: a systematic review and meta-analysis of observational studies. *JAMDA*. 2017;18:451.e1-.e11.
42. Roncero C, Campuzano AI, Quintano JA, Molina J, Perez J, Miravittles M. Cognitive status among patients with chronic obstructive pulmonary disease. *International Journal of COPD*. 2016;11:543-51.
43. Cleutjens FAHM, Spruit MA, Ponds RWHM, Vanfleteren LEGW, Franssen F, Gijzen C, *et al*. Cognitive impairment and clinical characteristics in patients with chronic obstructive pulmonary disease. *Chronic Respiratory Disease*. 2018;15(2):91-102.
44. Pierobon A, Bottelli ES, Ranzini L, Bruschi C, Maestri R, Bertolotti G, *et al*. COPD patients' self-reported adherence, psychosocial factors and mild cognitive impairment in pulmonary rehabilitation. *International Journal of COPD*. 2017;12:2059-67.
45. Meek P, Lareau SC, Anderson D. Memory for symptoms in COPD patients: how accurate are their reports? *European Respiratory Journal*. 2001;18(3):474-81.
46. Baird C, Lovell J, Johnson M, Shiell K, Ibrahim JE. The impact of cognitive impairment on self-management in chronic obstructive pulmonary disease: a systematic review. *Respiratory Medicine*. 2017;129:130-9.
47. Zogg JB, Woods SP, Saucedo JA, Wiebe JS, Simoni JM. The role of prospective memory in medication adherence: a review of an emerging literature. *Journal of Behavioral Medicine*. 2012;35:47-62.
48. Chang SS, Chen S, McAvay GJ, Tinetti ME. Effect of coexisting chronic obstructive pulmonary disease and cognitive impairment on health outcomes in older adults. *Journal of the American Geriatric Society*. 2012;60(10):1839-46.
49. Dassen FCM, Houben K, Van Breukelen GJP, Jansen A. Gamified working memory training in overweight individuals reduces food intake but not body weight. *Appetite*. 2018;124:89-98.
50. Houben K, Wiers RW, Jansen A. Getting a grip on drinking behavior: training working memory to reduce alcohol abuse. *Psychological Science*. 2011;22(7):968-75.
51. Dodd JW, Charlton RA, Van den Broek MD, Jones PW. Cognitive dysfunction in patients hospitalized with acute exacerbation of COPD. *Chest*. 2013;144(1):119-27.
52. Yazar EE, Aydin S, Gunluoglu G, Kamat S, Gungen AC, Yildiz P. Clinical effects of cognitive impairment in patients with chronic obstructive pulmonary disease. *Chronic Respiratory Disease*. 2018;15(3):306-14.
53. Schure MB, Borson S, Nguyen HQ, Trittschuh EH, Thielke SM, Pike KC, *et al*. Associations of cognition with physical functioning and health-related quality of life among COPD patients. *Respiratory Medicine*. 2016;114:46-52.

54. Vancampfort D, Koyanagi A, Ward PB, Veronese N, Carvalho AF, Solmi M, *et al.* Perceived stress and its relationship with chronic medical conditions and multimorbidity among 229,293 community-dwelling adults in 44 low- and middle-income countries. *American Journal of Epidemiology*. 2017;186(8):979-89.
55. Roiland RA, Lin F, Phelan C, Chapman BP. Stress regulation as a link between executive function and pre-frailty in older adults. *Journal of Nutrition, Health and Aging*. 2015;19(8):828-38.
56. Shields GS, Moons WG, Slavich GM. Better executive function under stress mitigates the effects of recent life stress exposure in young adults. *Stress*. 2017;20(1):75-85.
57. Esser RW, Stoeckel MC, Kirsten A, Watz H, Taube K, Lehmann K, *et al.* Structural brain changes in patients with COPD. *Chest*. 2016;149(2):426-34.
58. Savage CC, Dixey PHA, Pennington C, Dodd JW. Visual rating assessment of cerebral atrophy and its relationship with cognitive function in chronic obstructive pulmonary disease. *BMJ Open Respiratory Research*. 2018;5(1):e000310.
59. Chen J, Lin I-T, Zhang H, Lin J, Zheng S, Fan M, *et al.* Reduced cortical thickness, surface area in patients with chronic obstructive pulmonary disease: a surface-based morphometry and neuropsychological study. *Brain Imaging and Behavior*. 2016;10:464-76.
60. Franklin TB, Saab BJ, Mansuy IM. Neural mechanisms of stress resilience and vulnerability. *Neuron*. 2012;75(6):747-61.
61. McEwen BS, Bowles NP, Gray JD, Hill MN, Hunter RG, Karatsoreos IN, *et al.* Mechanisms of stress in the brain. *Nature Neuroscience*. 2015;18:1353-63.
62. Arnsten AFT. Stress weakens prefrontal networks: molecular insults to higher cognition. *Nature Neuroscience*. 2015;18(10):1376-85.
63. Lupien SJ, Juster R-P, Raymond C, Marin M-F. The effects of chronic stress on the human brain: from neurotoxicity, to vulnerability, to opportunity. *Frontiers in Neuroendocrinology*. 2018;49:91-105.
64. Datta D, Arnsten AFT. Loss of prefrontal cortical higher cognition with uncontrollable stress: molecular mechanisms, changes with age, and relevance to treatment. *Brain Sciences*. 2019;9:113.
65. Schram MT, Euser SM, De Craen AJ, Witteman JC, Frölich M, Hofman A, *et al.* Systemic markers of inflammation and cognitive decline in old age. *Journal of the American Geriatric Society*. 2007;55(5):708-16.
66. Crisan AF, Oancea C, Timar B, Fira-Mladinescu O, Crisan A, Tudorache V. Cognitive impairment in chronic obstructive pulmonary disease. *PLoS One*. 2014;9(7):e102468.
67. Tulek B, Atalay NB, Yildirim G, Kanat F, Suerdem M. Cognitive function in chronic obstructive pulmonary disease: relationship to global initiative for chronic obstructive lung disease 2011 categories. *Respirology*. 2014;19(6):873-80.

68. Zhang X, Cai X, Shi X, Zheng Z, Zhang A, Guo J, *et al.* Chronic obstructive pulmonary disease as a risk factor for cognitive dysfunction: a meta-analysis of current studies. *Journal of Alzheimer's disease*. 2016;52:101-11.
69. Ortapamuk H, Naldoken S. Brain perfusion abnormalities in chronic obstructive pulmonary disease: comparison with cognitive impairment. *Annals of Nuclear Medicine*. 2006;20(2):99-106.
70. Antonelli Incalzi R, Marra C, Giordano A, Calcagni ML, Cappa A, Basso S, *et al.* Cognitive impairment in chronic obstructive pulmonary disease - a neuropsychological and SPECT study. *Journal of Neurology*. 2003;250(3):325-32.
71. Andreou G, Vlachos F, Mekanikas K. Effects of chronic obstructive pulmonary disease and obstructive sleep apnea on cognitive functions: evidence for a common nature. *Sleep Disorders*. 2014;2014:768210.
72. Olaithe M, Bucks RS, Hillman DR, Eastwood PR. Cognitive deficits in obstructive sleep apnea: insights from a meta-review and comparison with deficits observed in COPD, insomnia, and sleep deprivation. *Sleep Medicine Reviews*. 2018;38:39-49.
73. Aras YG, Tunc A, Gungen BD, Gungen AC, Aydemir Y, Demiyurek BE. The effects of depression, anxiety and sleep disturbances on cognitive impairment in patients with chronic obstructive pulmonary disease. *Cognitive Neurodynamics*. 2017;11(6):565-71.
74. Dal Negro RW, Bonadiman L, Tognella S, Bricolo FP, Turco P. Extent and prevalence of cognitive dysfunction in chronic obstructive pulmonary disease, chronic non-obstructive bronchitis, and in asymptomatic smokers, compared to normal reference values. *International Journal of COPD*. 2014;26(9):675-83.
75. Blondell SJ, Hammersley-Mather R, Veerman JL. Does physical activity prevent cognitive decline and dementia? A systematic review and meta-analysis of longitudinal studies. *BMC Public Health*. 2014;14:510.
76. Kobayashi Y, Takahashi Y, Seki T, Kaneta T, Amarume K, Kasai M, *et al.* Decreased physical activity associated with executive dysfunction correlates with cognitive impairment among older adults in the community: a retrospective analysis from the Kurihara project. *Dementia and Geriatric Cognitive Disorders Extra*. 2016;6(2):350-60.
77. Falck RS, Davis JC, Liu-Ambrose T. What is the association between sedentary behaviour and cognitive function? A systematic review. *British Journal of Sports Medicine*. 2017;51(10):800-11.
78. Diet and nutrition in dementia and cognitive decline. London, England: Academic Press; 2015.
79. Cavalieri M, Ropele S, Petrovic K, Pluta-Fuerst A, Homayoon N, Enzinger C, *et al.* Metabolic syndrome, brain magnetic resonance imaging, and cognition. *Diabetes Care*. 2010;33(12):2489-95.

80. Falkowski J, Atchison T, DeButte-Smith M, Weiner MF, O'Bryant S. Executive functioning and the metabolic syndrome: a Project FRONTIER study. *Archives of Clinical Neuropsychology*. 2014;29:47-53.
81. Peng T-C, Chen W-L, Wu L-W, Chang Y-W, Kao T-W. Sarcopenia and cognitive impairment: a systematic review and meta-analysis. *Clinical Nutrition*. 2019;S0261-5614(19):33200-5.
82. Cabett Cipolli G, Sandes Yassuda M, Aprahamian I. Sarcopenia is associated with cognitive impairment in older adults: a systematic review and meta-analysis. *Journal of Nutrition, Health and Aging*. 2019;23:525-31.
83. Moon JH, Moon JH, Kim KM, Choi SH, Lim S, Park KS, *et al.* Sarcopenia as a predictor of future cognitive impairment in older adults. *Journal of Nutrition, Health and Aging*. 2016;20(5):496-502.
84. Pascual-Leone A, Amedi A, Fregni F, Merabet LB. The plastic human brain cortex. *Annual Review of Neuroscience*. 2005;28:377-401.
85. Pascual-Leone A, Freitas C, Oberman L, Horvath JC, Halko M, Bashir S, *et al.* Characterizing brain cortical plasticity and network dynamics across the age-span in health and disease with TMS-EEG and TMS-fMRI. *Brain Topography*. 2011;24:302.
86. Antonelli Incalzi R, Corsonello A, Trojano L, Pedone C, Acanfora D, Spada A, *et al.* Cognitive training is ineffective in hypoxemic COPD: a six-month randomized controlled trial. *Rejuvenation Research*. 2008;11:239-50.
87. Conway ARA, Kane MJ, Engle RW. Working memory capacity and its relation to general intelligence. *Trends in Cognitive Sciences*. 2003;7(12):547-52.
88. Barnett SM, Ceci SJ. When and where do we apply what we learn? A taxonomy for far transfer. *Psychological Bulletin*. 2002;128(4):612-37.
89. Burke SN, Barnes CA. Neural plasticity in the ageing brain. *Nature Reviews Neuroscience*. 2006;7:30-40.
90. Fan BJY, Wong RYM. Effect of cognitive training on daily function in older people without major neurocognitive disorder: a systematic review. *Geriatrics*. 2019;4(3):44.
91. Zhang H, Huntley J, Bhome R, Holmes B, Cahill J, Gould RL, *et al.* Effect of computerised cognitive training on cognitive outcomes in mild cognitive impairment: a systematic review and meta-analysis. *BMJ Open*. 2019;9:e027062.
92. Teixeira-Santos AC, Moreira CS, Magalhaes R, Magalhaes C, Pereira DR, Leite J, *et al.* Reviewing working memory training gains in healthy older adults: a meta-analytic review of transfer for cognitive outcomes. *Neuroscience and Biobehavioral Reviews*. 2019;103:163-77.
93. Wason PC, Evans JSBT. Dual processes in reasoning? *Cognition*. 1974;3(2):141-54.
94. Petty RE, Cacioppo JT. The elaboration likelihood model of persuasion. *Advances in Experimental Social Psychology*. 1986;19(1):123-205.

95. Kahneman D. A perspective on judgment and choice: mapping bounded rationality. *American Psychologist*. 2003;58(9):697-720.
96. Kahneman D. *Thinking, fast and slow*. New York, NY: Farrar, Straus and Giroux; 2011.
97. Evans JSBT, Stanovich KE. Dual-process theories of higher cognition: advancing the debate. *Perspectives on Psychological Science*. 2013;8(3):223-41.
98. Kantorowski A, Wan ES, Homsy D, Kadri R, Richardson CR, Moy ML. Determinants and outcomes of change in physical activity in COPD. *ERJ Open Research*. 2018;4(3):00054-2018.
99. Clarenbach CF, Sievi N, Haile SR, Brack T, Brutsche MH, Frey M, et al. Determinants of annual change in physical activity in COPD. *Respirology*. 2017;22(6):1133-9.
100. Tudor-Locke C, Craig CL, Aoyagi Y, Bell RC, Croteau KA, De Bourdeaudhuij I, et al. How many steps/day are enough? For older adults and special populations. *International Journal of Behavioral Nutrition and Physical Activity*. 2011;8:80.
101. Van de Bool C, Mattijssen-Verdonschot C, Van Melick PPMJ, Spruit MA, Franssen FME, Wouters EFM, et al. Quality of dietary intake in relation to body composition in patients with chronic obstructive pulmonary disease eligible for pulmonary rehabilitation. *European Journal of Clinical Nutrition*. 2014;68(2):159-65.
102. Laudisio A, Costanzo L, Di Gioia C, Delussu AS, Traballese M, Gemma A, et al. Dietary intake of elderly outpatients with chronic obstructive pulmonary disease. *Archives of Gerontology and Geriatrics*. 2016;64:75-81.
103. Chambaneau A, Filatre M, Jubert L, Bremond M, Filatre E. Nutritional intake, physical activity and quality of life in COPD patients. *International Journal of Sports Medicine*. 2016;37:730-7.







# Chapter 2

Clinical outcome and cost-effectiveness of a 1-year nutritional intervention programme in COPD patients with low muscle mass: the randomized controlled NUTRAIN trial

Martijn van Beers

Maureen P. M. H. Rutten-van Mölken

Coby van de Bool

Melinde Boland

Stef P. J. Kremers

Frits M. E. Franssen

Ardy van Helvoort

Harry R. Gosker

Emiel F. Wouters

Annemie M. W. J. Schols

*Clinical Nutrition. 2020;39(2):405-13*

## Abstract

**Background and aims:** The efficacy of nutritional intervention to enhance short- and long-term outcomes of pulmonary rehabilitation in COPD is still unclear, hence this paper aims to investigate the clinical outcome and cost-effectiveness of a 12-month nutritional intervention strategy in muscle-wasted COPD patients.

**Methods:** Prior to a 4-month pulmonary rehabilitation programme, 81 muscle-wasted COPD patients (51% males, aged  $62.5 \pm 0.9$  years) with moderate airflow obstruction ( $FEV_1$   $55.1 \pm 2.2\%$  predicted) and impaired exercise capacity ( $W_{max}$   $63.5 \pm 2.4\%$  predicted) were randomized to 3 portions of nutritional supplementation per day (enriched with leucine, vitamin D and polyunsaturated fatty acids) [NUTRITION] or PLACEBO (phase 1). In the unblinded 8-month maintenance phase (phase 2), both groups received structured feedback on their physical activity level assessed by accelerometry. NUTRITION additionally received 1 portion of supplemental nutrition per day and motivational interviewing-based nutritional counselling. A 3-month follow-up (phase 3) was included.

**Results:** After 12 months, physical capacity measured by quadriceps muscle strength and cycle endurance time were not different, but physical activity was higher in NUTRITION than in PLACEBO ( $\Delta$  1030 steps/day,  $p=0.025$ ). Plasma levels of the enriched nutrients ( $p<0.001$ ) were higher in NUTRITION than PLACEBO. Trends towards weight gain in NUTRITION and weight loss in PLACEBO led to a significant between-group difference after 12 months ( $\Delta$  1.54 kg,  $p=0.041$ ). The HADS anxiety and depression scores improved in NUTRITION only ( $\Delta$  -1.92 points,  $p=0.037$ ). Generic quality of life (EQ-5D) was decreased in PLACEBO but not in NUTRITION (between-group difference after 15 months 0.072 points,  $p=0.009$ ). Overall motivation towards exercising and healthy eating was high and did not change significantly after 12 months; only amotivation towards healthy eating yielded a significant between-group difference ( $\Delta$  1.022 points,  $p=0.015$ ). The cost per quality-adjusted life-year after 15 months was €16,750.

**Conclusions:** Nutritional intervention in muscle-wasted patients with moderate COPD does not enhance long-term outcome of exercise training on physical capacity but ameliorates plasma levels of the supplemented nutrients, total body weight, physical activity and generic health status, at an acceptable increase of costs for patients with high disease burden.

## Introduction

Extra-pulmonary pathology enhances disease burden in patients with chronic obstructive pulmonary disease (COPD) (1). Decreased muscle mass is associated with increased prevalence of osteoporosis, lower muscle strength and exercise performance, and increased mortality risk (2). Furthermore, COPD is characterized by impaired muscle oxidative metabolism, which cannot simply be explained by disuse alone (3, 4) and may predominate in patients with low muscle mass (5).

Exercise training is a cornerstone of pulmonary rehabilitation (PR) (6), and improves lower limb muscle strength, exercise performance, and in some studies also daily physical activity level (7, 8). Protein and specific nutrient supplementation could potentiate the effects of PR or represent an alternative to exercise training in severely deconditioned patients. In contrast to other nutrients including creatine and nitrate (9), to date only polyunsaturated fatty acids (PUFA) (10) and vitamin D (11) have been effective as single-nutrient add-on treatment in COPD by enhancing inspiratory muscle strength (11) and maximal oxygen uptake (10, 11) during PR which could be via immune modulation and/or boosting oxidative metabolism. Multimodal nutritional interventions are interesting to explore because the regulation of muscle mass and metabolism is controlled by multiple tightly intertwined pathways (12). In the NUTRAIN trial (13), we developed a multimodal stepwise nutritional intervention approach including high-quality protein enriched with leucine, vitamin D, and omega-3 PUFA, based on evidence regarding the mode of action of these nutrients on skeletal muscle maintenance, combined with reported deficiencies in COPD. The nutritional supplementation was supplemented by nutritional counselling. This approach was investigated in COPD patients with low muscle mass as a susceptible group for accelerated functional decline (14). The results of the first phase, which took place during PR, have been reported earlier (13). During this phase nutritional supplementation improved nutritional status but did not enhance lower limb muscle strength or muscle mass regain. A remarkable between-group difference was shown in physical activity level, largely due to a decline in the PLACEBO group only, whereas the NUTRITION group remained stable, indicating that this nutritional strategy might be effective to attenuate the decline in physical activity typically observed in patients with COPD (15). Exercise-induced muscle fatigue is a commonly reported symptom in COPD (16), and nutritional modulation thereof or nutritional effects beyond the muscle (i.e., the brain or cardiovascular system) (17, 18) might explain this observation. Accordingly, Calder *et al.* (19) showed in a 3-month randomized controlled trial that supplementation with high-quality protein enriched with omega-3 PUFA and

vitamin D, reduced walking exercise-induced fatigue (measured using the Borg scale) in COPD patients with low muscle mass.

We hypothesized that improved nutritional status and positive effects on daily physical activity level might also enhance the long-term efficacy of the intervention with quadriceps muscle strength (QMS) as primary outcome. The current paper aims to investigate the clinical outcome of the overall nutritional management strategy of the NUTRAIN trial, including phase 1, the 8-month maintenance phase after completion of PR (phase 2) and the 3-month follow-up (phase 3). Furthermore, the cost-effectiveness of the intervention after 15 months is explored.

## Methods

### *Study design*

The trial was integrated in the outpatient PR program of 7 hospitals in The Netherlands (Sint Anna hospital in Geldrop, Maxima Medical Centre in Veldhoven [until March 2013], Laurentius hospital in Roermond, Sint Jans Gasthuis hospital in Weert, Elkerliek hospital in Helmond, Maastricht University Medical Centre and Catharina hospital in Eindhoven), supervised by CIRO, a center of expertise for patients with chronic organ failure in Horn. Participants were recruited during the assessment period at CIRO when eligible for outpatient PR at one of the hospitals. Patients with severe COPD referred for inpatient PR at CIRO were not included. Every participant underwent an interview with a chest physician at admission; the study physician checked eligibility of potential participants during this visit and asked participants whether they could be contacted by the researcher. The study was registered at [clinicaltrials.gov](https://clinicaltrials.gov) (NCT01344135) and the medical ethics committee of Maastricht University Medical Centre granted ethical approval (NL34927068.10/MEC 11-3-004). All participants provided written informed consent.

The intervention was divided into a 4-month nutritional intervention (phase 1) and an 8-month maintenance phase (phase 2). In short, in the double-blind controlled intervention phase 1, 81 patients were randomized to the PLACEBO ( $n=39$ ) or NUTRITION group ( $n=42$ ). Both groups underwent a supervised center-based exercise training program and were advised to consume 3 oral nutritional supplements daily. The NUTRITION product provided protein, carbohydrates, fat and micronutrients, and was enriched with leucine, omega-3 PUFA and vitamin D (Nutricia NV, Zoetermeer, The Netherlands). The PLACEBO product did not

comprise the active components, but consisted of a flavored non-caloric cloudified aqueous solution. For details see Van de Bool *et al.* (13).

In the open-label maintenance phase (phase 2), both groups received feedback on their daily physical activity behavior twice, based on accelerometry. This was done in order to offer both groups a follow-up intervention aimed at maintaining long-term efficacy of the exercise training. In addition, participants in NUTRITION were advised to take 1 oral nutritional supplement per day and received four nutritional counselling sessions. These were provided by trained nurses, and aimed to optimize participants' dietary habits and maximize adherence to a healthy diet and the supplement regime. The counselling was based on self-determination theory (SDT) (20), which assumes that autonomous forms of motivation are essential to achieve lasting behavioral change, and it was operationalized through motivational interviewing (MI) (21). After the 8-month maintenance phase, participants were followed up for three months, without any intervention taking place in either group (phase 3). Figure 1 depicts a flowchart of the study; Figure 2 outlines the study design.

### *Patients*

Patients with COPD (post-bronchodilator  $FEV_1/FVC < 0.7$ ) referred for outpatient PR between September 2011 and April 2014 were eligible for participation if they had low muscle mass, defined as a fat-free mass index (FFMI) below the sex- and age-specific 25th percentile FFMI values (22). Exclusion criteria were age under 18 years old, allergy or intolerance to components of the study product, investigator's uncertainty about patients' willingness or ability to comply with the protocol requirements, inability to stop current supplement use, participation in any other study involving investigational or marketed products concomitantly or less than two weeks prior to entry into the study, pregnancy, or life-threatening diseases.

### *Outcomes*

Quadriceps muscle strength (QMS), assessed by dynamometry (System 4 Pro; Biodex Medical Systems Inc., New York, USA) served as the primary outcome of this study. QMS, body composition measured by dual energy x-ray absorptiometry (DEXA) (Lunar Prodigy system; GE Healthcare, Madison, WI, USA), cycle endurance time (CET) on a cycle ergometer (Carefusion, Houten, The Netherlands) (determined using the constant work rate cycling endurance test at 75% of the peak workload), and fasting plasma levels of vitamin D, eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA) and leucine were measured at baseline (before




Figure 1

Consort flow chart






Figure 2  
Study design

entering PR), and after phase 1 and 2. Habitual dietary intake of the last month was assessed at these time points by trained dietitians using a validated cross-check dietary history method and calculated using the Dutch Food Composition Database. Mental and physical health status, assessed by the Hospital Anxiety and Depression Scale (HADS) (23), EuroQoL Five-Dimensions Questionnaire (EQ-5D-3L) (24), St George's Respiratory Questionnaire (SGRQ) (25) and physical activity level (PAL) as assessed by daily steps by a tri-axial GT3X Actigraph accelerometer (Health One Technology, Fort Walton Beach, FL, USA) were measured at baseline and after phase 1, 2 and 3. Post-bronchodilator forced expiratory volume in 1 second (FEV<sub>1</sub>) and forced vital capacity (FVC) were assessed by standardized equipment (Masterlab®; Jaeger, Würzburg, Germany), in accordance with the latest GOLD guidelines (26). The Dutch value set was used to transform EQ-5D scores into utilities (27). Adopting such a wide range of outcome measures is in line with the recommendations of the joint ATS/ERS task force on outcome measures in COPD to use a multi-outcome approach (28).

Participants' motivational profiles towards exercising and healthy eating were assessed using the Behavioral Regulation in Exercise Questionnaire-2 (BREQ-2) (29) and the Regulation of Eating Behaviors Scale (REBS) (30), respectively, at baseline and after phase 1, 2 and 3. These questionnaires are based on SDT and they provide six subscale scores (see (20) for details). Answers were given on a 5-point Likert scale in both questionnaires. Scores on the BREQ range from 1 (lowest possible motivation) to 5 (highest possible motivation). Scores on the REBS range from 0 (lowest possible motivation) to 10 (highest possible motivation). Internal consistency measured by Cronbach's  $\alpha$  was adequate for all scales, except for identified regulation towards exercising ( $\alpha = 0.319$  after phase 2). Consequently, this scale was excluded from subsequent analyses.

## Costs

Total costs (not only related to COPD) were calculated from a healthcare and a societal perspective. The healthcare perspective included all costs covered by the healthcare budget. The societal perspective additionally included travel costs and costs of productivity loss due to absence from paid work (absenteeism). Healthcare utilization, travel distance and absenteeism were assessed in a resource use questionnaire after phase 1, 2 and 3.

The intervention costs included the reported number of supervised exercise training sessions and their duration, the number of nutritional supplements used, the number of educational counselling sessions and their duration, the number

of consultations for physical activity level feedback, the number of nutritional counselling sessions and the travel distance to health care providers.

Standard unit costs were determined using Dutch guideline prices (31) and inflated to 2015 values using the general consumer price index (32). The costs of medication were obtained from the Drug Information System of the Dutch National Health Care Institute and included value added tax and pharmacist dispensing fees (33). The productivity costs were estimated using the Friction Cost Approach, which assumes that productivity loss occurs as long as a sick employee is not replaced (the friction period) (34). We used a friction period of 85 days. The unit costs are shown in Supplementary Tables 1 through 3.

### *Statistical analyses*

Statistical analyses were performed using Stata, in accordance with the intention-to-treat principle.

Plasma levels of the supplemented nutrients, body composition, motivational profile and physical function were analyzed using a linear repeated measurement model with correlated error terms and unstructured covariance matrix. The model included treatment, time (i.e., the measurements at 4 and 12 months) and treatment by time interactions. Data from patients who discontinued the trial prematurely were included in the analyses up to the point of drop-out.

Costs were also analyzed in a linear repeated measures model with correlated error terms and unstructured covariance matrix. The dependent variable was total costs in a certain phase, and the explanatory variables were time (4, 12 and 15 months) and the interaction of treatment and time. The results were used to predict the mean costs per treatment group for each phase. Total costs were calculated as the sum of the predicted costs of the rehabilitation phase, maintenance phase and follow-up phase.

To assess differences in health outcomes between the two treatment groups, we also used linear repeated measurement models with correlated errors and unstructured covariance matrix. Explanatory variables were time (4, 12 [and, if applicable, 15] months) and the interaction of treatment and time. We calculated the number of quality-adjusted life-years (QALYs) for each patient as the area under the predicted utility curve, using linear interpolation between two measurements.

The costs per gained QALY were calculated as the difference in total costs between the two groups divided by the difference in QALYs. Uncertainty around this cost-effectiveness ratio was estimated by performing 5000 bootstrap replications.

### *Power calculation*

The power calculation was based on the INTERCOM study assuming a 10% between-group difference in peak torque assuming maintenance of skeletal muscle strength in NUTRITION during follow-up and a standard deviation of 5 Nm in peak torque. Allowing for 25% drop-out yielded a subgroup size of  $n=60$ . Patient inclusion was prematurely discontinued because the test product could not be produced within the appropriate quality specifications due to discontinuation of the supply of one of the ingredients, but the sample size was justifiable based on other RCTs published in the meantime (13).

## **Results**

### *Patients*

Baseline characteristics did not differ between the groups, except for a higher peak workload in NUTRITION. The study population consisted of 51% males, aged  $62.5 \pm 0.9$  years, and was characterized by low diffusion capacity (DLCO  $49.4 \pm 1.7\%$ ), moderate airflow limitation ( $FEV_1$   $55.1 \pm 2.2\%$  predicted), normal to low BMI ( $22.7 \pm 0.3$  kg/m<sup>2</sup>), impaired exercise capacity ( $W_{max}$   $63.5 \pm 2.4\%$  predicted), and low FFMI ( $15.8 \pm 1.6$  kg/m<sup>2</sup>) (see Table 1).

### *Physical functioning*

No between-group difference was observed in improved physical capacity (QMS and CET) (see Table 2). When QMS was normalized for appendicular lean mass or fat-free mass, the between-group difference was not significant either (data not shown). Physical activity level was significantly lower compared to baseline in PLACEBO whereas it remained stable in NUTRITION. This led to a significant between-group difference on number of steps per day ( $p=0.025$ ) (see Figure 3). Activity intensity did not differ significantly over time or between the groups: in both groups across all time points, patients spent around 70% in sedentary physical activity (PA), 23% in lifestyle PA, 6% in light PA and 1% in moderate-to-vigorous PA.

Table 1

## Baseline characteristics

	<b>NUTRITION</b>	<b>PLACEBO</b>
<i>General</i>		
Gender (% male)	42.9%	59.0%
Age (years)	62.8±1.3	62.2±1.3
Current smokers (%)	19.0%	31.6%
Low SES (no or only primary education (%))	5.6%	13.3%
Employed (%)	33.3%	19.4%
Number of exacerbations in the last year	1.1±0.2	1.4±0.3
<i>Lung function</i>		
FEV <sub>1</sub> (% predicted)	57.0±3.3	53.0±2.8
FVC (% predicted)	102.8±2.9	100.6±2.7
FEV <sub>1</sub> /FVC (%)	44.4±2.0	41.6±1.8
Functional residual capacity (% predicted)	139.1±4.3	146.7±5.9
Residual volume (% predicted)	143.4±6.3	153.1±8.5
DLCO (% predicted)	51.4±2.2	47.1±2.5
<i>Physical function</i>		
Quadriceps muscle strength (Nm)	121.5±6.4	118.0±6.6
Cycle endurance time (s)	319.0±35.2	231.5±12.0
Peak workload (W <sub>max</sub> )	84.6±5.2	72.5±3.8
Peak workload (% predicted)	69.5±3.3**	57.0±3.4
Physical activity level (steps per day)	4716.7±327.2	4516.6±379.3
<i>Dietary intake</i>		
Energy (kcal)	2426.0±168.0	2442.8±174.2
Protein (g)	91.1±5.5	89.9±5.5
Carbohydrates (g)	263.1±19.9	266.8±17.5
Fat (g)	103.1±9.8	105.4±12.3
Polyunsaturated fatty acids (g)	20.2±2.0	21.5±2.7
Cholesterol (mg)	251.5±25.2	239.1±25.2
Dietary fiber (g)	21.9±1.3	21.7±1.6

Table 1 (continued)

	<b>NUTRITION</b>	<b>PLACEBO</b>
Calcium (mg)	1083.5±96.1	977.3±67.5
Vitamin A (µg)	1241.4±96.5	1192.5±145.8
Vitamin C (mg)	84.1±6.1	85.9±6.9
Vitamin D (µg)	5.3±0.4	5.1±0.6
Vitamin E (mg)	14.6±1.3	13.9±1.2
<i>Supplemented plasma nutrient levels</i>		
Leucine (µmol/l)	147.7±3.6	153.5±3.5
Vitamin D (nmol/l)	54.0±4.5	45.3±3.1
Eicosapentaenoic acid (mg/l)	1.1±0.1	1.2±0.1
Docosahexaenoic acid (mg/l)	2.9±0.2	3.2±0.2
<i>Body composition</i>		
Total weight (kg)	64.3±1.6	65.0±1.7
Bone mineral content (g)	2331.3±73.0	2414.7±82.3
Appendicular skeletal muscle mass (kg)	17.4±0.6	18.4±0.6
Fat mass (kg)	20.0±1.0	19.0±1.3
Lean mass (kg)	42.0±1.3	43.6±1.2
Body mass index (kg/m <sup>2</sup> )	22.9±0.4	22.6±0.5
Fat-free mass index (kg/m <sup>2</sup> )	15.7±0.3	15.9±0.2
<i>Patient-reported outcomes</i>		
EuroQoL 5-dimensions questionnaire	0.8±0.0	0.8±0.0
St George's Respiratory Questionnaire	49.9±2.4	50.1±2.6
Hospital Anxiety and Depression Scale - total	10.9±1.2	10.2±1.1

All data given as mean ± standard error, unless noted otherwise. DLCO: diffusion capacity of the lung for carbon monoxide; FEV<sub>1</sub>: forced expiratory volume in 1 second; FVC: forced vital capacity; Nm: Newtonmeter; SES: socio-economic status. \*\**p*<0.01.




Figure 3

Development of physical activity level as change from baseline. # within-group  $p < 0.05$  (compared to baseline); \* between-group  $p < 0.05$ .

The change in physical activity level from baseline to T12 correlated significantly with the change in appendicular skeletal muscle mass index (ASMI) ( $r = 0.371$ ,  $p = 0.022$ ) but not that in total body weight ( $r = 0.088$ , n.s.) in the entire study population.

### Dietary intake and plasma nutrient status

Overall dietary habits represented a typical Western diet and no significant within- or between-group changes were identified throughout the trial on any macronutrient or micronutrient, except for cholesterol intake (see Supplementary Table 4). Cholesterol intake was significantly higher in PLACEBO after 12 months

Table 2

Within- and between-group differences in plasma nutrient status, physical function, body composition and patient-reported outcomes, after 12 months

	Within-group				Between-group	
	NUTRITION		PLACEBO		adj. M (SE)	Z
Measure	adj. M (SE)	Z	adj. M (SE)	Z		
Physical function						
QMS (Nm)	10.35±3.51	2.95**	10.05±3.46	2.90**	0.30±4.92	0.06
CET (s)	107.1±62.9	1.70*	200.3±64.5	3.11**	-93.2±89.8	1.04
PAL (steps/day)	358.8±339.9	1.06	-671.3±330.3	-2.03*	1030.1±459.8	2.24*
Supplemented plasma nutrient levels						
Leucine (μmol/l)	10.41±5.57	1.87	-6.08±5.80	-1.05	16.49±8.01	2.06*
Vitamin D (nmol/l)	15.10±2.69	5.61***	1.25±2.77	0.45	13.85±3.84	3.60***
EPA (mg/l)	5.42±1.66	3.27**	-3.34±1.64	-2.04*	8.76±1.95	4.49***
DHA (mg/l)	7.26±2.69	2.70**	-4.69±2.67	-1.76	11.97±3.47	3.45**
Body composition						
Total weight (kg)	0.64±0.53	1.21	-0.90±0.54	-1.67	1.54±0.76	2.04*
BMC (g)	-9.96±13.81	-0.72	-22.21±14.12	-1.57	12.26±19.73	0.62
ASM (kg)	0.17±0.21	0.83	-0.22±0.21	-1.02	0.39±0.29	1.32
Fat mass (kg)	0.86±0.50	1.71	-0.47±0.51	-0.91	1.33±0.72	1.86



Table 2 (continued)

Measure	Within-group				Between-group	
	NUTRITION		PLACEBO		adj. M (SE)	Z
	adj. M (SE)	Z	adj. M (SE)	Z		
Patient-reported outcomes						
EQ-5D	-0.003±0.03	-0.10	-0.07±0.03	-2.63**	0.07±0.04	1.86
SGRQ	1.43±1.89	0.75	-0.71±1.97	-0.04	2.14±2.73	0.78
HADS - total	-1.92±0.92	-2.09*	-1.50±0.97	-1.54	-0.42±1.30	-0.32

Values shown as change from baseline. ASM: appendicular skeletal muscle mass; BMC: bone mineral content; CET: cycle endurance time; DHA: docosahexaenoic acid; EPA: eicosapentaenoic acid; EQ-5D: EuroQoL 5-dimensions questionnaire; HADS: Hospital Anxiety and Depression Scale; Nm: Newtonmeter; PAL: physical activity level; QMS: quadriceps muscle strength; SGRQ: St George's Respiratory Questionnaire. \*  $p<0.05$ ; \*\*  $p<0.01$ ; \*\*\*  $p<0.001$ .




Figure 4

Change in total body weight from baseline to T12. \* between-group  $p < 0.05$ .

compared to baseline ( $p = 0.001$ ), leading to a significant between-group difference ( $p = 0.003$ ). At baseline, 31.2% of participants had vitamin D insufficiency, whereas 57.5% had vitamin D deficiency.

Plasma vitamin D, EPA and DHA levels were significantly increased compared to baseline in NUTRITION. EPA levels were significantly decreased compared to baseline in PLACEBO. Leucine, vitamin D, EPA and DHA levels significantly differed between groups (see Table 2).

### Body composition

Total body weight, muscle mass and fat mass tended to decrease within PLACEBO and tended to increase within NUTRITION from baseline to the end of phase 2. This resulted in a significant between-group difference in total body weight change, in favor of NUTRITION ( $\Delta 1.54 \pm 0.76$  kg,  $p = 0.041$ ) (see Table 2 and Figure 4).

Table 3

Within- and between-group differences in physical activity level and patient-reported outcomes, after 15 months.

Measure	Within-group				Between-group	
	NUTRITION		PLACEBO		adj. M (SE)	Z
PAL (steps per day)	adj. M (SE)	Z	adj. M (SE)	Z	796.1±496.9	1.60
	-38.1±355.5	-0.11	-834.2±362.8	-2.30*		
EQ-5D	0.012±0.02	0.49	-0.060±0.02	-2.53*	0.072±0.03	2.12*
SGRQ	0.16±2.11	0.08	2.72±2.19	1.24	-2.56±3.00	-0.86
HADS - total	-0.85±0.97	-0.87	-1.74±0.98	-1.78	0.89±1.38	0.65

Values shown as change from baseline. EQ-5D: EuroQoL 5-dimensions questionnaire; HADS: Hospital Anxiety and Depression Scale; PAL: physical activity level; SGRQ: St George's Respiratory Questionnaire. \*  $p<0.05$ .

## *Patient-reported outcomes*

Total HADS score significantly improved in NUTRITION ( $p=0.037$ ). In PLACEBO, the EQ-5D score decreased significantly ( $p=0.009$ ) (see Table 2). The SGRQ did not differ significantly within or between groups. After phase 3, EQ-5D utility was still significantly decreased compared to baseline in PLACEBO ( $p=0.011$ ), whereas the between-group difference was also significant in favor of NUTRITION ( $p=0.034$ ) (see Table 3).

Interestingly, the change in EQ-5D scores from baseline to T12 correlated significantly with the change in total body weight ( $r=0.406$ ,  $p=0.008$ ) and ASMI ( $r=0.456$ ,  $p=0.003$ ) in the entire study population.

## *Motivational profile*

Motivation towards both exercising and healthy eating was already high before the counselling sessions started. The mean scores for intrinsic and identified motivation towards exercising before the counselling started were 4.14 and 4.03 out of 5, respectively. These figures were 7.69 and 8.42 out of 10 for intrinsic and identified motivation towards healthy eating. In NUTRITION, amotivation towards exercising decreased ( $p=0.015$ ) while in PLACEBO identified motivation towards healthy eating decreased ( $p=0.019$ ). Only amotivation towards healthy eating differed significantly between groups ( $p=0.015$ ) (see Table 4).

## *Intervention costs*

Intervention costs are presented in Supplementary Table 1. From a healthcare perspective, the mean intervention costs of the total intervention were €2,233 per patient in NUTRITION and €1,372 in PLACEBO. From a societal perspective, these costs were €2,265 and €1,404, respectively. The between-group difference is mainly due to the costs of the nutritional supplements. Supervised exercise training was the main driver of the intervention costs in both groups.

## *Total costs and cost-effectiveness*

Tables 5 and 6 show the cost-effectiveness analysis after phase 2 and 3. The differences in various categories of healthcare utilization over 12 and 15 months are presented in Supplementary Tables 2 and 3. After phase 2, the costs in NUTRITION were estimated to be €1,529 (certainty 87%) higher than in PLACEBO from the healthcare perspective and €2,829 (certainty 98%) higher from the societal perspective. After phase 3, costs in NUTRITION were €670 (certainty 64%)

Table 4

Within- and between-group differences in motivational profile, after 12 months

Measure	Within-group				Between-group	
	NUTRITION		PLACEBO		adj. M (SE)	Z
	adj. M (SE)	Z	adj. M (SE)	Z		
Exercise						
Intrinsic	0.069±0.11	0.60	0.011±0.12	0.09	0.058±0.16	0.36
Introjected	-0.183±0.16	-1.12	0.212±0.17	1.25	-0.396±0.23	-1.72
External	0.006±0.10	0.06	-0.006±0.10	-0.06	0.012±0.13	0.09
Amotivation	-0.232±0.10	-2.43*	-0.011±0.10	-0.17	-0.215±0.13	-1.71
Healthy eating						
Intrinsic	-0.094±0.26	-0.36	0.060±0.27	0.22	-0.154±0.37	-0.42
Integrated	0.040±0.29	0.14	-0.230±0.31	-0.76	0.271±0.40	0.67
Identified	-0.028±0.24	-0.12	-0.577±0.24	-2.35*	0.549±0.34	1.60
Introjected	-0.437±0.29	-1.50	-0.644±0.30	-0.21	-0.373±0.41	-0.91
External	-0.297±0.38	-0.78	-0.122±0.39	-0.31	-0.175±0.53	-0.33
Amotivation	-0.492±0.30	-1.44	0.593±0.31	1.92	-1.022±0.42	-2.43*

Values shown as change from baseline. \*  $p<0.05$ .

Table 5

Cost-effectiveness and cost-utility: 12-month time horizon

	<b>NUTRITION</b>	<b>PLACEBO</b>	<b>Difference (95% UI)</b>
<b>Costs (€)</b>			
Healthcare perspective	8,094	6,565	1,529 (-1,007 - 4,343)
Societal perspective	9,404	6,575	2,829 (100 - 5,802)
<b>Effects</b>			
QALYs	0.66	0.64	0.025 (-0.02 - 0.07)
PAL (steps/day)	4,960	3,940	1,020 (17 - 1,990)
Percentage of patients with +4 units improvement in SGRQ	0.36	0.43	-0.07 (-0.38 - 0.22)
<b>ICER (€)</b>	<b>HCP</b>	<b>SCP</b>	
Costs / QALY gained	61,160	113,160	
Costs / 1000 additional steps	1,499	2,774	
Costs / additional patient with +4 units improvement in SGRQ	dominated	dominated	

\*  $p < 0.05$ . HCP: healthcare perspective; PAL: physical activity level; QALY: quality-adjusted life year; SCP: societal perspective; SGRQ: St George's Respiratory Questionnaire; UI: uncertainty interval.

higher from the healthcare perspective and €2,401 (certainty 92%) higher from the societal perspective.

Both from a healthcare and societal perspective, the point-estimates of costs pointed towards higher costs in NUTRITION and a higher physical activity level in NUTRITION after phase 2, and a higher number of QALYs in NUTRITION after phase 3. The probability that the nutritional intervention was cost-effective at a willingness-to-pay of €20,000 and €80,000 per QALY was 27% and 59% after phase 2 and 55% and 82% after phase 3 (data not shown).

## Discussion

This paper investigated the clinical benefits and cost-efficacy of a 12-month multimodal, stepwise nutritional intervention strategy in patients with moderate

Table 6

Cost-effectiveness and cost-utility: 15-month time horizon

	<b>NUTRITION</b>	<b>PLACEBO</b>	<b>Difference (95% UI)</b>
<b>Costs (€)</b>			
Healthcare perspective	9,548	8,878	670 (-2,468 - 4,001)
Societal perspective	11,324	8,923	2,401 (-854 - 5,824)
<b>Effects</b>			
QALYs	0.86	0.82	0.04 (-0.01 - 0.10)
PAL (steps/day)	4,576	3,780	814 (-335 - 1,838)
Percentage of patients with +4 units improvement in SGRQ	0.35	0.23	0.12 (-0.16 - 0.39)
<b>ICER (€)</b>	<b>HCP</b>	<b>SCP</b>	
Costs / QALY gained	16,750	60,025	
Costs / 1000 additional steps	823	2,950	
Costs / additional patient with +4 units improvement in SGRQ	5,583	20,008	

\*  $p < 0.05$ . HCP: healthcare perspective; PAL: physical activity level; QALY: quality-adjusted life year; SCP: societal perspective; SGRQ: St George's Respiratory Questionnaire; UI: uncertainty interval.

COPD and low muscle mass. Physical capacity, as primary outcome, was improved in both groups after 12 months, but NUTRITION took 1,030 additional steps per day compared to PLACEBO. This difference exceeds the minimal important difference to reduce risk of hospital admission (35), and it cannot be explained by greater energy substrate availability, as total macronutrient intake was not significantly different between the groups at T12 (data not shown). This indicates that nutrient-specific mechanisms could explain the dissociation between changes in QMS and physical activity via various synergistic pathways including immune modulation (PUFA, vitamin D) (36, 37), enhancing muscle oxidative metabolism (PUFA) (38) and modifying muscle maintenance regulation (PUFA, leucine) (39, 40).

Plasma levels of the supplemented nutrients after 12 months were indeed significantly higher in NUTRITION compared to both baseline and PLACEBO, indicating that the phased intervention strategy was feasible and overall compliance was good.

2 Trends towards weight gain in NUTRITION and weight loss in PLACEBO after 12 months led to a significant between-group difference in total body weight. A tendency to lose weight in PLACEBO was also found after 12 months in a trial focusing on protein-calorie supplementation by Weekes *et al.* (41). Changes in muscle and fat mass paralleled total weight change but were not significant, possibly due to lack of power. Remarkably, in contrast to weight change, the subtle change in muscle mass correlated significantly with the change in PA.

Previous studies in weight-losing and muscle-wasted patients with advanced COPD showed positive short-term (1-4 months) effects of protein-calorie supplementation as adjunct to exercise training on physical capacity and muscle mass (42-44), but limited evidence is available in less advanced airflow obstruction. The INTERCOM trial, which investigated a combined nutritional and exercise intervention in the subgroup of patients with low muscle mass and reduced cycle exercise capacity, also showed positive effects on muscle mass and physical capacity in less advanced COPD patients with low muscle mass (14). However, it could not disentangle the relative contributions of the exercise and nutritional interventions to physical performance, because the muscle-wasted control group only received usual care. In the 20-month maintenance phase of the INTERCOM trial, physical capacity remained improved in the subgroup of treated muscle-wasted patients, but 6-minute walking distance declined below baseline in those receiving usual care (14). Whereas the nutritional intervention strategy had no short- or long-term added value on physical capacity in the current trial (13), there were sustained positive effects on body weight, plasma nutrient levels and physical activity and a trend towards improved mental health, which could explain the difference in generic health status after follow-up.

Both EQ-5D and physical activity levels remained stable in NUTRITION and declined in PLACEBO after 12 and 15 months. These parallel changes could mean that generic health status is influenced by daily physical activity level *via* modulation of physiological and / or mental health.

A qualitative study in a random sample of 22 NUTRAIN participants after phase 2 identified perceived competence and autonomous motivation as determinants of a physically active lifestyle (45). The limited between-group motivational differences could be partly explained by high levels of both intrinsic and extrinsic motivation and low levels of amotivation already apparent before counselling. Admission to PR already requires being motivated and participating in an extensive study such as this one even more so. Even so, slight changes in motivational profile may indicate some effect of the nutritional counselling.



The incremental cost-effectiveness ratio (ICER) of €61,160 after phase 2 decreased to €16,750 after phase 3, because of a higher gain in QALYs and lower difference in costs after phase 3. Whether an intervention is cost-effective or not depends on the threshold value of the willingness-to-pay for a QALY. In The Netherlands this threshold depends on disease burden (defined as proportional shortfall) (46). A higher threshold applies for conditions with a higher disease burden. There are three categories, defining a low burden (proportional shortfall of 0.1-0.4 QALYs), medium (0.41-0.7) and high burden of disease (0.71-1), corresponding of threshold values of €20,000, €50,000 and €80,000 per QALY, respectively (47). Hence, an ICER below €20,000 (after phase 3) is generally considered to be cost-effective. With an ICER of around €60,000 after phase 2, the current nutritional intervention would be cost-effective for COPD patients with a disease burden of at least 0.7 QALYs. However, this estimation is associated with high uncertainty illustrated by a small proportion of bootstrap replications below a threshold value of €80,000 after phase 2.

A weakness of the current study includes the unblinding after PR, which was done because it was considered unethical to continue placebo supplementation during the maintenance phase. Secondly, as already mentioned, the current study included quite a small number of patients, in line with many other comparable studies. Generalizability of the study is limited by the fact that only patients referred for PR were included. Thirdly, not achieving the sample size specified beforehand could limit our ability to detect significant results. Finally, the counselling sessions were given by nurses who lacked detailed knowledge on the domains to be improved. More detailed and tailored advice could improve adherence to a healthy lifestyle and increase health gains.

## Conclusion

A stepwise multimodal nutritional intervention strategy, consisting of targeted nutritional supplementation and nutritional counselling, in muscle-wasted patients with moderate COPD does not enhance long-term outcome of exercise training on physical capacity but ameliorates plasma levels of the supplemented nutrients, body weight, physical activity and generic health status. This intervention increases total health care costs to a degree that might be considered acceptable for patients with a high disease burden.

## References

1. Barnes PJ. Chronic obstructive pulmonary disease: effects beyond the lungs. *PLoS Medicine*. 2010;7(3):e1000220.
2. Maltais F, Decramer M, Casaburi R, Barreiro E, Burelle Y, Debigare R, *et al*. An official American Thoracic Society / European Respiratory Society statement: update on limb muscle dysfunction in chronic obstructive pulmonary disease. *American Journal of Respiratory and Critical Care Medicine*. 2014;189:e15-e62.
3. Konokhova Y, Spendiff S, Jagoe RT, Aare S, Kapchinsky S, MacMillan NJ, *et al*. Failed upregulation of TFAM protein and mitochondrial DNA in oxidatively deficient fibers of chronic obstructive pulmonary disease locomotor muscle. *Skelet Muscle*. 2016;6:10.
4. Gifford JR, Trinity JD, Kwon O-S, Layec G, Garten RS, Park S-Y, *et al*. Altered skeletal muscle mitochondrial phenotype in COPD: disease versus disuse. *Journal of Applied Physiology*. 2018;124:1045-53.
5. Van de Bool C, Gosker HR, Van den Borst B, Op den Kamp CM, Slot IG, Schols AM. Muscle quality is more impaired in sarcopenic patients with chronic obstructive pulmonary disease. *JAMDA*. 2016;17:415-20.
6. Spruit MA, Singh SJ, Garvey C, ZuWallack R, Nici L, Rochester C, *et al*. An official American Thoracic Society / European Respiratory Society statement: Key concepts and advances in pulmonary rehabilitation. *American Journal of Respiratory and Critical Care Medicine*. 2013;188(8):e13-e64.
7. De Sousa Pinto JM, Martin-Nogueras AM, Calvo-Arenillas JI, Ramos-Gonzalez J. Clinical benefits of home-based pulmonary rehabilitation in patients with chronic obstructive pulmonary disease. *Journal of Cardiopulmonary Rehabilitation and Prevention*. 2014;34:355-9.
8. Pleguezuelos E, Perez ME, Guirao L, Samitier B, Ortega P, Vila X, *et al*. Improving physical activity in patients with COPD with urban walking circuits. *Respiratory Medicine*. 2013;107:1948-56.
9. Shepherd AI, Wilkerson DP, Dobson L, Kelly J, Winyard PG, Jones AM, *et al*. The effect of dietary nitrate supplementation on the oxygen cost of cycling, walking performance and resting blood pressure in individuals with chronic obstructive pulmonary disease: a double blind placebo controlled, randomised control trial. *Nitric Oxide*. 2015;48.
10. Broekhuizen R, Wouters EF, Creutzberg EC, Weling-Scheepers CA, Schols AM. Polyunsaturated fatty acids improve exercise capacity in chronic obstructive pulmonary disease. *Thorax*. 2005;60:376-82.
11. Hornikx M, Van Remoortel H, Lehouck A, Mathieu C, Maes K, Gayan-Ramirez G, *et al*. Vitamin D supplementation during rehabilitation in COPD: a secondary analysis of a randomized trial. *Respiratory Research*. 2012;13:84.

12. Remels AH, Gosker HR, Langen RC, Schols AMWJ. The mechanisms of cachexia underlying muscle dysfunction in COPD. *Journal of Applied Physiology*. 2013;114(9):1253-62.
13. Van de Bool C, Rutten EPA, Van Helvoort A, Franssen FM, Wouters EF, Schols AM. A randomized clinical trial investigating the efficacy of targeted nutrition as adjunct to exercise training in COPD. *Journal of Cachexia, Sarcopenia and Muscle*. 2017;8:748-58.
14. Van Wetering CR, Hoogendoorn M, Broekhuizen R, Geraerts-Keeris GJW, De Munck DRAJ, Rutten-Van Molken MPMH. Efficacy and costs of nutritional rehabilitation in muscle-wasted patients with chronic obstructive pulmonary disease in a community-based setting: a prespecified subgroup analysis of the INTERCOM trial. *JAMDA*. 2010;11:179-87.
15. Troosters T, Sciurba F, Battaglia S, Langer D, Valluri SR, Martino L, et al. Physical inactivity in patients with COPD: a controlled multi-center pilot-study. *Respiratory Medicine*. 2010;104:1005-11.
16. Amann M, Regan MS, Kobitarty M, Eldridge MW, Boutellier U, Pegelow DF, et al. Impact of pulmonary system limitations on locomotor muscle fatigue in patients with COPD. *American Journal of Physiology*. 2010;299:R314-24.
17. Robinson LE, Mazurak VC. N-3 polyunsaturated fatty acids: relationship to inflammation in healthy adults and adults exhibiting features of metabolic syndrome. *Lipids*. 2013;48:319-32.
18. Janssens W, Lehouck A, Carremans C, Bouillon R, Mathieu C, Decramer M. Vitamin D beyond bones in chronic obstructive pulmonary disease: time to act. *American Journal of Respiratory and Critical Care Medicine*. 2009;179:630-6.
19. Calder PC, Laviano A, Lonnqvist F, Muscaritoli M, Ohlander M, Schols AM. Targeted medical nutrition for cachexia in COPD: a randomized, controlled trial. *Journal of Cachexia, Sarcopenia and Muscle*. 2018;9:28-40.
20. Deci EL, Ryan RM. *Intrinsic motivation and self-determination in human behavior*. New York, NY: Plenum Press; 1985.
21. Resnicow K, Dilorio C, Soet JE, Ernst D, Borrelli B, Hecht J. Motivational interviewing in health promotion: it sounds like something is changing. *Health Psychology*. 2002;21:444-51.
22. Schutz Y, Kyle UUG, Pichard C. Fat-free mass index and fat mass index percentiles in Caucasians aged 18-98y. *International Journal of Obesity and Related Metabolic Disorders*. 2002;26:953-60.
23. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatrica Scandinavica*. 1983;67:361-70.
24. EuroQol Group. EuroQol - a new facility for the measurement of health-related quality of life. *Health Policy*. 1990;16:199-208.

25. Jones PW, Quirk FH, Baveystock CM. The St George's Respiratory Questionnaire. *Respiratory Medicine*. 1991;85(Supplement B):25-31.
26. Vestbo J, Hurd SS, Agusti AG, Jones PW, Vogelmeier C, Anzueto A, *et al*. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *American Journal of Respiratory and Critical Care Medicine*. 2013;187(4):347-65.
27. Lamers LM, McDonnell J, Stalmeier PF, Krabbe PF, Busschbach JJ. The Dutch tariff: results and arguments for an effective design for national EQ-5D valuation studies. *Health Economics*. 2006;15:1121-32.
28. Cazzola M, MacNee W, Martinez FJ, Rabe KF, Franciosi LG, Barnes PJ, *et al*. Outcomes for COPD pharmacological trials: from lung function to biomarkers. *European Respiratory Journal*. 2008;31:416-69.
29. Markland D, Tobin VJ. A modification to the Behavioral Regulation in Exercise Questionnaire to include an assessment of amotivation. *Journal of Sport and Exercise Psychology*. 2004;26:191-6.
30. Pelletier LG, Dion SC, Slovinec-D'Angelo M, Reid R. Why do you regulate what you eat? Relationships between forms of regulation, eating behaviors, sustained dietary behavior change and psychological adjustment. *Motivation and Emotion*. 2004;28(3):245-77.
31. Hakkaart-Van Roijen L, Tan S, Bouwmans C. Handleiding voor kostenonderzoek, methoden en standaard kostprijzen voor economische evaluaties in de gezondheidszorg. College voor Zorgverzekeringen; 2010.
32. CBS Statline. Consumentenprijzen: inflatie vanaf 1963-2013. Available from: <http://statline.cbs.nl/StatWeb/publication/?VW=T&DM=SLNL&PA=70936NED&D1=0&D2=%281-34%29-I&HD=081020-1258&HDR=T&STB=G1>.
33. College of Health Insurance (CVZ). GIP databank. Available from: <http://www.gipdatabank.nl>.
34. Koopmanschap MA, Rutten FF, Van Ineveld BM, Van Roijen L. The friction cost method for measuring indirect costs of disease. *Journal of Health Economics*. 1995;14:171-89.
35. Demeyer H, Burtin C, Hornikx M, Camillo CA, Van Remoortel H, Langer D, *et al*. The minimal important difference in physical activity in patients with COPD. *PLoS One*. 2016;11:e0154587.
36. Giudetti AM, Cagnazzo R. Beneficial effects of n-3 PUFA on chronic airway inflammatory diseases. *Prostaglandins and Other Lipid Mediators*. 2012;99:57-67.
37. Solidoro P, Bellocchia M, Facchini F. The immunobiological and clinical role of vitamin D in obstructive lung diseases. *Minerva Medica*. 2016;107(Supplement 1-3):12-9.
38. Chen PB, Yang J, S., Park Y. Adaptations of skeletal muscle mitochondria to obesity, exercise, and polyunsaturated fatty acids. *Lipids*. 2018;53:271-8.

39. Nicastro H, Artioli GG, Dos Santos Costa A, Solis MY, Ribeiro da Luz C, Blachier F, *et al.* An overview of the therapeutic effects of leucine supplementation on skeletal muscle under atrophic conditions. *Amino Acids*. 2011;40:287-300.
40. Rondanelli M, Faliva M, Monteferrario F, Peroni G, Repaci E, Allieri F, *et al.* Novel insights on nutrient management of sarcopenia in elderly. *BioMed Research International*. 2015:524948.
41. Weekes CE, Emery PW, Elia M. Dietary counselling and food fortification in stable COPD: a randomized controlled trial. *Thorax*. 2009;64:326-31.
42. Sugawara K, Takahashi H, Kasai C, Kiyokawa N, Watanabe T, Fujii S, *et al.* Effects of nutritional supplementation combined with low-intensity exercise in malnourished patients with COPD. *Respiratory Medicine*. 2010;104:1883-9.
43. Laviolette L, Lands LC, Daultbaev N, Saey D, Milot J, Provencher S, *et al.* Combined effect of dietary supplementation with pressurized whey and exercise training in chronic obstructive pulmonary disease: a randomized, controlled, double-blind pilot study. *Journal of Medicinal Food*. 2010;13:589-98.
44. Gurgun A, Deniz S, Argin M, Karapolat H. Effects of nutritional supplementation combined with conventional pulmonary rehabilitation in muscle-wasted chronic obstructive pulmonary disease: a prospective, randomized and controlled study. *Respirology*. 2013;18:495-500.
45. Stewart KF, Meis JJM, Van de Bool C, Janssen DJA, Kremers SPJ, Schols AMWJ. Maintenance of a physically active lifestyle after pulmonary rehabilitation in patients with COPD: a qualitative study toward motivational factors. *JAMDA*. 2014;15(9):655-64.
46. Van de Wetering EJ, Stolk EA, Van Exel NJ, Brouwer WB. Balancing equity and efficiency in the Dutch basic benefits package using the principle of proportional shortfall. *European Journal of Health Economics*. 2013;14:107-15.
47. Kosteneffectiviteit in de praktijk. Diemen, The Netherlands: Zorginstituut Nederland; 2015.

Appendix

Supplementary Table 1

Intervention costs (in euros, 2015)

		NUTRITION				PLACEBO			
	Unit cost	Source	Mean cost (all patients)	Any use (%)	Mean use	Mean cost (users)	Any use (%)	Mean use	Mean cost (users)
Phase 1									
Supervised exercise training sessions	33	a	1,235±24	100	36	1,197±42	95	39	1,279±17
Reference priced product, # supplements	2.63	b	293±39	100	215	565±43	-	-	-
Nutrition placebo, # supplements	-	-		-	-	-	95	272	-
Education, training sessions	8.93	a	125		14	125	100	14	125
Travel expenses, public transport/car, per km	0.19	a	31±1	100	160	30±1	100	171	32±0.5
Phase 2									
Feedback on physical activity, consultations	33	a	37±3	69	1.2	39±4	59	1.1	34±5
Nutritional counselling, consultations	17	a		55	3.5	59±2	-	-	-
Reference priced product, # supplements	2.63	b		50	208	548±35	-	-	-

Supplementary Table 1 (continued)

				NUTRITION			PLACEBO		
	Unit cost	Source	Mean cost (all patients)	Any use (%)	Mean use	Mean cost (users)	Any use (%)	Mean use	Mean cost (users)
Travel expenses, public transport/car, per km	0.19	a	0.9±0.1	74	7	1.2±0.2	59	2.3	0.4±0.06
Total costs from health-care perspective			1,818±76			2,233±104			1,372±50
Total costs from societal perspective			1,850±77			2,265±105			1,404±51

Costs are reported in euros and expressed as mean ± standard deviation. Sources of unit costs used in the analysis: (a) Dutch guidelines for pharmacoeconomic research, (b) Z-index.

Supplementary Table 2

Unit costs, data sources, mean unadjusted use of resources and associated costs over 12 months, as reported by the patients (excluding costs related to the intervention program)

		NUTRITION				PLACEBO		
	Unit cost	Source	Any use (%)	Mean use	Mean cost	Any use (%)	Mean use	Mean cost
Costs from healthcare perspective								
General practitioner	17-50	a	79	4.60	144±19	74	4.54	138±35
Office consultations	33	a	69	3.00	98±15	67	2.38	79±20
Home visits	50	a	7	0.17	8±5	8	0.44	22±13
Phone contacts	17	a	17	0.33	6±2	21	0.59	10±4
Practice nurse consultations	17	a	17	0.40	7±3	26	0.77	13±4
Specialist, visits	91	a	76	5.40	492±98	72	6.85	623±173
Emergency department, visits	166	a	14	0.52	136±70	13	0.26	66±41
Physiotherapist, visits	33	a	48	21.4	706±151	36	18.9	624±165
Dietician, visits	30	a	38	0.86	26±8	41	0.95	32±10
Podiatrist, visits	34	b	24	1.10	37±12	13	0.58	20±10
Speech therapist, visits	30	a	2	0.05	1±1	5	0.15	5±3
Occupational therapy, visits	33	a	33	0.88	34±12	28	0.44	14±4
Home care, hours	22-78		10	2.66	74±44	15	9.57	305±148
Hospital stay, days			19	2.31	1,541±687	18	0.79	378±161



Supplementary Table 2 (continued)

				NUTRITION			PLACEBO		
	Unit cost	Source	Any use (%)	Mean use	Mean cost	Any use (%)	Mean use	Mean cost	
No intensive care unit, days	476	a	19	2.03	965±433	18	0.79	378±161	
Intensive care unit, days	2,015	a	10	0.29	576±324	0	0	0	
Medication									
Drugs for obstructive airway diseases	-	c	86	-	1,085±195	79	-	1,308±324	
Drugs for cardiovascular disease	-	c	40	-	171±48	49	-	188±47	
Drugs for diabetes	-	c	5	-	31±29	8	-	5±3	
Costs from societal perspective									
Travel expenses, public transport/car, per km	0.19	a	83	179	34±7	82	157	30±8	
Productivity loss, absenteeism hours	32-38	a	10	27	1,303±666	0	0	0	

Costs are reported in euros and expressed as mean ± standard deviation. Sources of unit costs used in the analysis: (a) Dutch guidelines for pharmacoeconomic research, (b) Z-index.

Supplementary Table 3

Unit costs, data sources, mean unadjusted use of resources and associated costs over 15 months, as reported by the patients (excluding costs related to the intervention program)

			NUTRITION				PLACEBO		
	Unit cost	Source	Any use (%)	Mean use	Mean cost	Any use (%)	Mean use	Mean cost	
Costs from healthcare perspective									
General practitioner	17-50	a	81	5.47	170±23	77	5.01	154±38	
Office consultations	33	a	71	3.50	116±17	67	2.56	85±20	
Home visits	50	a	12	0.21	11±6	10	0.51	26±15	
Phone contacts	17	a	17	0.40	7±3	21	0.64	11±5	
Practice nurse consultations	17	a	17	0.52	9±4	26	0.77	13±4	
Specialist, visits	91	a	81	6.10	555±102	74	7.79	709±182	
Emergency department, visits	166	a	21	0.60	154±70	23	0.36	93±42	
Physiotherapist, visits	33	a	50	27.6	910±198	51	26.4	870±212	
Dietician, visits	30	a	43	1.00	30±8	49	1.26	42±11	
Podiatrist, visits	34	b	24	1.21	41±13	13	0.66	22±11	
Speech therapist, visits	30	a	2	0.05	1±1	8	0.26	8±4	
Occupational therapy, visits	33	a	33	0.88	34±12	28	0.44	14±4	
Home care, hours	22-78		12	2.70	76±44	151	10.0	326±161	
Hospital stay, days			21	2.66	1,704±725	28	1.64	821±285	

Supplementary Table 3 (continued)

	Unit cost	Source	NUTRITION			PLACEBO		
			Any use (%)	Mean use	Mean cost	Any use (%)	Mean use	Mean cost
No intensive care unit, days	476	a	21	2.37	1,128±484	28	1.62	769±272
Intensive care unit, days	2,015	a	10	0.29	576±324	3	0.03	52±52
Medication								
Drugs for obstructive airway diseases	-	c	86	-	1,198±203	79	-	1,453±333
Drugs for cardiovascular disease	-	c	40	-	185±53	49	-	220±55
Drugs for diabetes	-	c	5	-	39±37	8	-	6±4
Costs from societal perspective								
Travel expenses, public transport/car, per km	0.19	a	86	216	41±8	82	210	40±9
Productivity loss, absenteeism hours	32-38	a	14	31.9	1,553±673	3	1.08	47±40

Costs are reported in euros and expressed as mean ± standard deviation. Sources of unit costs used in the analysis: (a) Dutch guidelines for pharmacoeconomic research, (b) Z-index.

Supplementary Table 4

Within- and between-group differences in dietary intake, after 12 months

Measure	Within-group				Between-group	
	NUTRITION		PLACEBO		adj. M (SE)	Z
Energy (kcal)	adj. M (SE)	Z	adj. M (SE)	Z	-178.20±153.30	-1.16
Protein (g)	-78.76±108.06	-0.73	99.44±111.07	0.90	-10.53±5.84	-1.80
Carbohydrate (g)	-4.45±4.19	-1.06	6.08±4.30	1.42	6.02±17.31	0.35
Fat (g)	4.29±12.13	0.35	-1.73±12.43	-0.14	-10.27±7.62	-1.35
PUFA (g)	-2.41±5.35	-0.45	7.86±5.52	1.42	0.09±1.67	0.05
Cholesterol (mg)	-0.77±1.19	-0.65	-0.86±1.23	-0.70	-67.48±22.48	-3.00**
Dietary fiber (g)	-13.68±16.47	-0.84	53.62±16.83	3.19**	0.14±1.44	0.10
Calcium (mg)	0.68±1.01	0.68	0.54±1.03	0.52	-135.29±94.04	-1.44
Vitamin A (µg)	-38.28±67.41	-0.57	97.01±68.80	1.41	-211.73±109.37	-1.94
Vitamin C (mg)	-56.46±87.17	-0.65	155.27±88.42	1.76	-7.41±12.27	-0.60
Vitamin D (µg)	3.93±9.00	0.44	11.34±9.26	1.22	-0.98±0.78	-1.26
Vitamin E (mg)	-0.82±0.58	-1.41	0.17±0.59	0.28	-0.45±1.64	-0.27
	-0.92±1.28	-0.72	-0.47±1.31	-0.36		

PUFA: polyunsaturated fatty acids. \*\*  $p<0.01$ .





# Chapter 3

Motivational regulations and lifestyle in patients with COPD and low muscle mass before and after pulmonary rehabilitation

EMBARGOED

Martijn van Beers

Jessie J. M. Meis

Coby van de Bool

Annemie M. W. J. Schols

Stef P. J. Kremers

*Submitted*

# Chapter 4

Cognitive impairment in chronic obstructive pulmonary disease: disease burden, determinants and possible future interventions

Martijn van Beers

Daisy J. A. Janssen

Harry R. Gosker

Annemie M. W. J. Schols



## Abstract

**Introduction:** Cognitive impairment (CI) is an important but under-recognized extra-pulmonary feature of chronic obstructive pulmonary disease (COPD). It is related to the burden of disability, worse health outcomes, and impaired self-management.

**Areas covered:** CI includes deterioration of a wide range of cognitive functions, such as memory and various executive functions. Risk of hospitalization might be higher in patients with COPD compared to those without, with CI negatively impacting the wellbeing of patients with COPD. Disease-specific factors such as hypoxemia and inflammation, lifestyle factors such as dietary insufficiencies and lack of physical activity, and comorbidities such as obstructive sleep apnea and depression are likely to synergistically contribute to the development of CI in COPD. Tailored interventions can possibly improve CI in COPD, but this needs further investigation.

**Expert commentary:** Further research is warranted involving the optimization of neuropsychological testing for screening and outcome assessment, longitudinal studies to investigate the development of CI in COPD over time, and randomized clinical trials to test the feasibility and efficacy of promising interventions.

## Introduction

Chronic obstructive pulmonary disease (COPD) is a chronic lung disease characterized by persistent respiratory symptoms and airflow limitation due to airway and / or alveolar abnormalities (1). It is well established that common extra-pulmonary comorbidities such as heart failure, osteoporosis and muscle wasting contribute significantly to the disease burden (2). Moreover, recent research shows that cognitive impairment (CI) is also a common and important, yet under-investigated comorbidity (3). This review will discuss various aspects related to CI in COPD: its prevalence and characteristics, related health outcomes, determinants, and possible interventions to maintain or improve cognitive functioning in patients with COPD.

## Cognitive functioning

Cognition can be defined as any brain function that enables an individual to perceive, register, store, retrieve and use information in order to adapt behavior to new situations and function in our environment (4). Cognition consists of many separate domains, including memory, working memory and attention (5).

Cognition is organized hierarchically and different cognitive functions can be classified as 'lower' or 'higher' (6) (see Figure 1). The lower functions form the basis for the higher functions, including the executive functions such as inhibition (choosing an option that is more rewarding in the longer term instead of the more immediately satisfying option when presented with a choice, for example refusing the attempt to smoke a cigarette) and cognitive flexibility (being able to swiftly shift mental resources, for example changing the type of physical activity undertaken in case of weather changes) (7). Executive functioning in turn underlies even more complex processes such as problem solving and decision making, for example when and how to undertake action in case of increased symptom severity (e.g., disease exacerbation or unintended weight loss).

Specific brain areas do not exclusively perform a single task, but many brain areas are specialized for certain types of tasks (see Figures 2 and 3). The language and memory systems are largely located in the temporal lobe, whereas the higher-order functions such as inhibition, cognitive flexibility, reasoning and decision-making are processed in the frontal lobe. Subcortical structures are located within the brain, including the hippocampus, which plays a central role in memory encoding; the limbic system, including the amygdala and cingulate cortex, which are involved




Figure 1

*A hierarchical view of cognitive processes. Adapted from Cleutjens et al. (4).*

in fear and pain processing; and the basal ganglia, including the caudate nucleus, which is involved in movement but also in learning and remembering (8).

Brain damage, for instance caused by brain atrophy or degeneration, can contribute to the development of CI. This is an inevitable part of aging, but acute and chronic disease can accelerate it. In the initial stages of CI development, impairments are noticeable, but not severe enough to impair activities of daily living; a stage called mild cognitive impairment (MCI) (5). MCI often, but not always, involves impairments in memory rather than in other domains (9), and it may progress into Alzheimer's disease (9) or remain stable (10).




Figure 2

View of the left cerebral hemisphere, indicating the areas mentioned in the text. Adapted from Patrick J. Lynch, medical illustrator, and Carl C. Jaffe, MD, cardiologist.

## Cognitive impairment in COPD

Patients with COPD display impairment in cognitive domains as diverse as attention, executive functioning (3, 11-13), (visual) memory and reproduction (14, 15), problem solving, concentration, logical and abstract reasoning, planning, coordination and organization (3). Thus, the pattern of impairments is diffuse, and not every patient is affected or suffers from CI to the same degree in different cognitive domains (16). This diffuse pattern resembles aging-related cognitive deterioration and is consistent with the view of CI in COPD as a manifestation of accelerated ageing (3). On the other hand, patients with Alzheimer's disease or other dementias show a much more specific pattern of CI.




Figure 3

*Sagittal view of the brain, indicating the areas mentioned in the text. Adapted from Patrick J. Lynch, medical illustrator, and C. Carl Jaffe, MD, cardiologist.*

## Prevalence

### Cross-sectional studies

A recent meta-analysis pooled 14 studies investigating the prevalence of CI in COPD patients (17). The average prevalence of any CI (5 studies,  $N=2,995$ ) was 32%. MCI was present in 25% of patients (11 studies,  $N=4,663$ ). Higher prevalence of CI was associated with respiratory disease severity, dependence in activities of daily living and poor quality of life (17).

Studies published after this meta-analysis (25 February 2016) show an even higher prevalence of CI (see Table 1). Two studies by Cleutjens *et al.* reported CI, determined by extensive neuropsychological assessment, in 56.7% and 41.5% of patients with COPD referred for pulmonary rehabilitation (PR) (18, 19). In the former

Table 1

Overview of studies investigating the prevalence of CI in COPD

Study	Country	Design	Setting	Population	Sample size	Gender (% male)	Age (M±SD)
Cleutjens, 2017a (18)	Netherlands	cross-sectional	PR	Clinically stable COPD admitted to PR	90	54.4	63.7±8.8
Cleutjens, 2017b (19)	Netherlands	cross-sectional	PR	Clinically stable COPD admitted to PR	76	60.5	62.7±8.7
				Non-CI controls with COPD	107	47.7	63.7±9.9
Cleutjens, 2017c (61)	Netherlands	cross-sectional	PR	Clinically stable COPD admitted to PR	157	50.3	62.9±9.4
Lopez-Torres, 2016 (25)	Spain	longitudinal	hospital	Patients hospitalized with and recovering from acute COPD exacerbation	62	75.8	68.3±7.4
Pierobon, 2017 (20)	Italy	observational, cross-sectional	PR	Stable (no exacerbation for the last 3 months) COPD patients with GOLD stage II-IV, group C-D	84	75.0	70.2±7.0
Park, 2018 (27)	USA	observational, longitudinal	NETT	Radiological evidence of bilateral emphysema, severe airflow obstruction and hyperinflation, and the ability to complete PR	307	59.6	66.2±5.7
Roncero, 2016 (21)	Spain	cross-sectional	respiratory medicine department	Ambulatory patients > 40 years with stable COPD	940	81.6	67.6±10.0
Samareh Fekri, 2017 (23)	Iran	cross-sectional	medical university	Patients with a history and symptoms of COPD	87	90.8	60.5±9.8
				Controls without COPD and CI	60	68.3	58.2±9.8

Table 1 (continued)

Study	Cognitive tests used	Criteria / cut-off for CI	% MCI	% ACI
Cleutjens, 2017a	Comprehensive neuropsychological test battery	Score less than 1SD below age-, gender- and education-specific mean of the MAAS study (24) on 2 subtests or more		56.7% (COPD) 13.3% (controls)
Cleutjens, 2017b	Comprehensive neuropsychological test battery	Score less than 1SD below age-, gender- and education-specific mean of the MAAS study (24) on 2 subtests or more		41.5%
Cleutjens, 2017c	Comprehensive neuropsychological test battery; MMSE	Score less than 1SD below age-, gender- and education-specific mean of the MAAS study on 2 subtests or more; MMSE < 24		5.7% (MMSE); 38.2% (test battery)
Lopez-Torres, 2016	MoCA	MoCA < 20		48.3% (exacerbating) 23.6% (discharge) 36.3% (stable)
Pierobon, 2017	MoCA; MMSE	MoCA performance within bottom 5% of the population		9.5%
Park, 2018	TMTB	Score more than 1.5SD above the normative mean of the current study (134.66s for those with < 12 years of education; 81.09s for those with less)		32.6%
Roncero, 2016	MMSE	MMSE < 27	39.4%	
Samareh Fekri, 2017	MMSE	Mild: MMSE 19-23 Moderate: MMSE 10-19 Severe: MMSE < 10	44.82% (COPD) 33.33% (controls)	6.89% (moderate CI, COPD) 3.33% (moderate CI, controls)

Table 1 (continued)

Study	Miscellaneous methodological remarks
Cleutjens, 2017a	Cognitive scores are split into different domains and composite scores are made
Cleutjens, 2017b	Cognitive scores are split into different domains and composite scores are made
Cleutjens, 2017c	Cognitive scores are split into different domains and composite scores are made
Lopez-Torres, 2016	Different numbers are given for gender and age distribution in the table compared with the figures. Different MoCA versions were used in order to eliminate practice effects; patients with dementia were excluded.
Pierobon, 2017	Patients with MMSE < 18.3 were excluded
Park, 2018	In- and exclusion criteria for the NETT trial did not specifically have cognitive research in mind: neurological disorders or medication which could affect cognition were not excluded, for instance
Roncero, 2016	
Samareh Fekri, 2017	Unconventional CI cut-offs; skewed gender distribution (91% men in the COPD group); many participants had an opium addiction or history of baking in traditional furnaces; non-smoking history was three times more common in the control group compared to the COPD group (36.7% versus 12.6%)

Studies included in the meta-analysis by Yohannes et al. (17) are not included in this table. CI: cognitive impairment; COPD: chronic obstructive pulmonary disease; GOLD: Global Initiative for Chronic Obstructive Lung Disease; MAAS: Maastricht Aging Study; MMSE: Mini-Mental State Examination; MoCA: Montreal Cognitive Assessment; NETT: National Emphysema Treatment Trial; PR: pulmonary rehabilitation; SD: standard deviation; TMTB: Trail Making Test, part B.



study only 13.3% of non-COPD age-matched controls suffered from CI. Pierobon *et al.* showed that the Montreal Cognitive Assessment (MoCA) score of 9.5% of patients admitted to PR was below the 5th percentile of a normal reference group, and only 38.1% scored higher than the median (20). This may be related to the fact that patients referred for PR generally have moderate to advanced disease. Roncero *et al.* showed CI in 39.4% of stable COPD patients. This study was large ( $N=940$ ), but it only assessed cognition using the Mini-Mental State Examination (MMSE) (21), which is useful as a screening tool but not as a diagnostic tool (22). Samareh Fekri *et al.* likewise administered the MMSE to 87 COPD patients and 60 age- and gender-matched individuals. MCI was present in 44.8% of patients with COPD and moderate CI in 6.9%, compared to 33.3% and 3.3%, respectively, in controls (23). It is not reported whether these percentages differ significantly, but the mean MMSE score in the COPD group was significantly lower than in the control group ( $22.51 \pm 2.4$  versus  $23.63 \pm 2.8$ , respectively).

Lopez-Torres *et al.* investigated cognition during hospitalization for an exacerbation, at discharge and after return to a stable state (25). Cognitive functioning measured by MoCA scores increased significantly from hospitalization to discharge and decreased again from discharge to stable state, but to a higher level than at hospitalization.

In summary, a wide range of prevalences is reported for CI in COPD. The heterogeneity of applied cognitive tests and cut-off points to define CI as well as diverse study populations (community-dwelling, pulmonary rehabilitation, hospitalized patients *et cetera*) likely contributes to this wide range.

## Longitudinal studies

Future research using longitudinal designs may elucidate the development of CI in COPD over several years and the influence of disease progression and other factors in different disease stages. The longitudinal studies already conducted have shown some interesting results. A self-reported diagnosis of COPD in midlife increased the risk of developing CI later in life (odds ratio [OR] 1.85, 95% confidence interval [CI95] 1.05-3.28), but COPD diagnosed later in life was non-significantly inversely related to CI (OR 0.30, CI95 0.08-1.24) (26). This surprising result might be explained by survival bias rather than the effects of COPD itself (26). In a recent study 32.6% of patients taking part in the National Emphysema Treatment Trial (NETT) were impaired on part B of the Trail Making Test (TMT) at baseline (27). This test measures task switching, which is part of executive functioning. No prevalence estimates were given for subsequent time points, but TMT-B performance was virtually unchanged over the 3-year follow-up. Most

remarkably, the sample could be distinguished into four clusters based on their trajectories of cognitive development over three years. One cluster (35.5%) had low baseline TMT-B scores and improving scores over time, the second (39.7%) had low baseline and worsening scores, the third (18.2%) had high baseline and worsening scores, and the fourth (6.5%) had high baseline and improving scores.

### *Cerebral abnormalities*

CI is associated with global and/or specific cerebral abnormalities, and many studies have found structural or functional abnormalities in patients with COPD, along with elevated serum levels of S100B, a putative marker for brain damage (28).

Cortical degeneration (29-32), increased occurrence of small vessel disease (33) or abnormal functional activation on a global level is uncommon in COPD, with a few exceptions. Two studies found overall increased white matter (WM) lesion volumes and decreased WM integrity (30, 31), and cortical thickness and volume were globally reduced in patients who were hospitalized for 30-45 days following an exacerbation (34).

Regional changes are more common. Gray matter (GM) was found to be decreased in many brain regions in both hemispheres, among others in the dorsolateral prefrontal cortex (34), which is involved in higher functions and working memory (35); different areas involved in visuospatial processing (36); the frontal cortex; and limbic and paralimbic structures (15, 29), which are mainly involved in emotion processing and memory (37). Disease duration was inversely related to GM volume in various, mainly subcortical, regions (12, 29), but not in others, including the hippocampus and amygdala (29).

Reduced WM integrity in the superior and middle frontal gyri and right occipital subcortical WM was shown in patients with moderate COPD (32). WM integrity was more reduced in the bilateral frontal subcortical areas, right temporal lobe and pons in severe compared to moderate COPD. Patients with an acute exacerbation who had been hospitalized for several weeks also showed reduced WM integrity compared to healthy controls in various (para)limbic regions (12). However, Cleutjens *et al.* (33) showed that cognitively weak and cognitively strong patients with COPD had equal amounts of WM hyperintensities (WMH).

Two studies investigated the relation between lung function and brain volume in healthy elderly. One study reported a significant correlation between forced expiratory volume in 1 second ( $FEV_1$ ) and overall brain atrophy and ventricle-to-brain ratio in men but not in women (36). The  $FEV_1$ /forced vital capacity (FVC)

ratio was correlated with WMH in the sample as a whole. Participants with and without chronic respiratory disease did not differ on any clinical or brain imaging parameter. Moreover, no control variables were included. The second study only showed a significant positive relationship between  $FEV_1$  and cerebellar WM volume, but no generalized cortical degeneration (39).

The hippocampus is interesting as it is vital in memory formation and learning, and only one of the two regions to display neurogenesis in the adult human brain (40). As such, decreased hippocampal volume could also indicate abnormalities in brain plasticity in patients with COPD, but findings are mixed. In one study, hippocampal volume was decreased in COPD compared to healthy controls and its size was related to partial oxygen pressure and oxygen saturation (41). Hippocampal volume did not differ between patients with mild-to-moderate COPD and those with severe COPD (41). In another study, however, hippocampal volume was not significantly different between patients with COPD and controls (42), and it has also been shown to not differ between cognitively strong and cognitively weak patients with COPD (33).

Functional abnormalities were reported in the left precentral and postcentral gyri and the left caudate nucleus when comparing patients with COPD with controls matched on age, sex and education (14). Evidence on resting state network and default mode network activity is mixed. These networks are distinct from others because their activity increases when the brain is not engaged with other tasks (43). Their increased activation could hinder cognition by interfering with task-related activity. One study found increased resting state network activity in patients with COPD compared to healthy controls, but this result was not significant anymore after controlling for oxygen saturation (14). Another study also found increased default network activation in patients with moderate COPD, but decreased activation in those with severe COPD (44). This result might reflect a compensatory response to damage inflicted by factors associated with COPD, and this response might be strongest in patients with moderate COPD (44).

The structural and functional abnormalities discussed above reflect a gradual influence of COPD on the brain. However, COPD can also heighten the risk of more acute events, such as stroke. This is likely due to high levels of systemic inflammation and oxidative stress, possibly as a result of smoking, leading to endothelial dysfunction, decreased vascular reactivity, thickening of the carotid artery wall and atherosclerotic plaque rupture (45). An elevated risk of stroke in patients with COPD was found in several studies (hazard ratios [HRs] 1.09 [CI95 0.91-1.31] and 1.24 [CI95 1.19-1.28]) (46, 47) and meta-analyses (HRs 1.30 [CI95 1.09-2.09]) (48, 49). The significance in the latter study was probably due to the

large sample size ( $N=132,017$  versus  $1,566$  in the former). HRs are roughly equal for the ischemic, hemorrhagic, intracerebral and subarachnoid hemorrhagic subtypes. However, in the weeks after an acute exacerbation, the HR increased to  $6.66$  (CI95  $2.42-18.20$ ) (46).

In conclusion, patients with COPD show an increased prevalence of many different types of cerebral abnormalities, compared to healthy controls. Patients with COPD have reduced gray and WM volume in various brain areas and WM integrity is compromised. It is still equivocal whether hippocampal volume is decreased in patients with COPD. There is evidence for abnormal functional activation, particularly in the resting state and default mode networks. The abnormalities are spread throughout the brain rather than concentrated in one or some areas. This could possibly explain the diffuse pattern of CI found in COPD.




Figure 4

*The possible relationships between cognitive impairment and clinical outcomes. Dotted lines represent conflicting literature concerning the relationship.*

## CI and health outcomes in COPD

CI has a wide range of adverse effects on self-management skills and other health outcomes in patients with COPD (see Figure 4).

### *Self-management skills*

Self-management skills are of major importance for patients with COPD. Symptoms may vary from day to day and increased symptoms may be caused by exacerbations. Patients need to recognize and act upon these exacerbations in an adequate manner (50). Moreover, patients need to adhere to their medication and need to adopt a healthy lifestyle, including a healthy diet and physical activity. Smokers need to quit smoking. This often requires a behavior change, which poses a demand on cognitive functioning (50). It is therefore feasible that CI limits the ability to cope with the daily challenges of living with COPD.

A recent systematic review revealed that there was just one study in which the impact of CI on self-management skills in patients with COPD was investigated (51). This particular study, including 100 participants with COPD, showed no relationship between cognitive functioning assessed with the MoCA and overall self-management abilities, as measured by the Self Management Ability Score 30 (SMAS30). SMAS30 assesses the following: taking initiatives, investment behavior, variety, multifunctionality, self-efficacy, and positive frame of mind. Moreover, living alone affected the interaction between cognitive functioning and self-management abilities (52). In fact, only among patients who lived alone, better cognitive functioning was related to lower self-management abilities (52). Emotional intelligence (defined as the capacity to understand and manage personal thoughts and feelings, as well as to positively influence interpersonal communication and social well-being), however, seems to be related to self-management abilities in COPD (53).

Meek *et al.* (54) showed a relationship between MMSE scores and the ability to accurately recall severity of fatigue and dyspnea in the previous two weeks. It is reasonable to assume that this might impact on the ability to recognize and act upon symptoms of an exacerbation, but the exact relationship remains unknown.

The relationship between CI and inhaler competency is well-known (51). An MMSE score of 23-24 points or less is predictive of poor inhaler technique, as are impaired executive functioning and impaired praxis. While some inhalers may be more difficult to use than others, recognizing CI is of major importance when prescribing inhaler devices as well as providing instructions to use them (51). Some studies suggest Turbhalers might be easier to use for patients

with CI than metered dose inhalers (51, 55). To our knowledge, however, there are no recommendations specifically on inhaler use for COPD patients with CI. Nevertheless, inhaler competency is a major consideration when prescribing inhalers and should be checked regularly.

Brega *et al.* (56) showed that older persons with impaired executive functioning were less likely to quit smoking than those with normal executive functioning. However, recent data did not confirm the relationship between executive cognitive dysfunction and smoking cessation (57). Moreover, another study even showed that persons over 75 years of age who quit smoking had lower cognitive functioning than persons who continued smoking (58). Differences in findings about the relationship between smoking and cognition might be explained by the fact that patients in the last study were older and different methods were used to assess cognitive functioning. Cleutjens *et al.* did not find a statistically significant difference in smoking status between patients with COPD entering PR with or without CI (19). Therefore, whether and to what extent CI limits the ability of patients with COPD to quit smoking remains unclear.

### *Health outcomes*

CI seems related to functional exercise capacity as measured by 6-minute walking distance (59, 60). Nevertheless, this was not confirmed among patients entering PR (19) or during an exacerbation (59).

Cleutjens *et al.* showed that the response after completion of a PR program was similar in patients with CI compared to those without (61). However, patients with CI were more likely to drop out compared to those without CI (23.3% versus 10.3%, respectively). Therefore, timely recognition of CI in patients entering PR seems paramount.

Several studies showed an inverse relationship between CI and HRQoL as assessed with the COPD Assessment Test (21, 60), the EuroQoL-5 dimensions questionnaire (21), or St George's Respiratory Questionnaire (22). Then again, other studies did not confirm this relationship (19, 62).

The systematic review of Baird *et al.* included four studies exploring the relationship between CI and disability and showed that CI is related to impairment in basic activities of daily living, instrumental activities of daily living, work and social activities (51). Martinez *et al.* found that COPD and CI have independent but additive effects on disability (63).

The co-existence of CI and COPD is associated with a more than fourfold increased risk of hospitalization for respiratory-related illnesses and a 34% higher risk of all-cause hospitalization compared to healthy controls, after controlling for various sociodemographic variables, smoking status and comorbidities (64). Moreover, CI seems to be related to an increased length of hospitalization (22). Data from the NETT trial did not show an association between impairment in executive functioning and frequency of hospitalization (65).

A study including stable patients with COPD even showed that drawing impairment predicted increased mortality risk (66). Nevertheless, Yohannes *et al.* found that MMSE scores did not predict 1-year all-cause mortality (67). Moreover, only a modest association was found between executive function and survival in the NETT trial (65).

In conclusion, CI may negatively impact self-management and health outcomes in COPD, but the current literature is conflicting and many questions remain. The conflicting literature may be explained by different methods used to assess cognitive functioning and criteria used to define CI. Moreover, there were major differences between the studied populations. The complex cause-effect relationships also make this field very challenging. COPD may influence self-management skills, but poor self-management in turn also worsens COPD disease progression, thereby creating a vicious cycle.

## **Determinants of COPD-induced cognitive impairment**

Factors such as hypoxemia, hypercapnia, inflammation and lifestyle factors may all contribute to structural and functional cerebral abnormalities and CI. Any individual factor probably does not explain a significant amount of CI, but their synergistic effects may be large (11, 68).

### *Hypoxemia and hypercapnia*

Hypoxemia and hypercapnia are hallmarks of severe COPD (69) which can also negatively affect cognition (70, 71). Hypoxemia alters the microenvironment around neurons (72) and induces impairments in spontaneous and task-stimulated neuronal activity (73, 74). Hypoxemia might underlie decreases in GM density (12, 15), and mild hypercapnia decreases functional connectivity in almost all brain lobes (75). The exact mechanisms through which this happens are still unclear (75). Diminished vasodilatation in response to hypoxemia or hypercapnia in patients with COPD might be one (76).

## *Inflammation*

Multiple lines of evidence indicate that systemic inflammation might underlie CI in COPD. Low-grade systemic inflammation is linked to decreased cognition in other conditions including obesity (77, 78) and metabolic syndrome (79, 80). In the general population c-reactive protein (CRP) and interleukin-6 levels were related to global cognition and executive functioning, whereas  $\alpha$ 1-antichymotrypsin was not (79). MoCA scores have been found to be negatively correlated with CRP, fibrinogen and erythrocyte sedimentation rate levels (82). The relationship between inflammation and cognition in stable COPD has not been thoroughly investigated yet. Next to low-grade systemic inflammation, patients with COPD may experience periods of an enhanced systemic inflammatory response related to disease exacerbations that will be discussed later.

## *Respiratory medication*

Long-acting  $\beta$ -adrenoreceptor agonists (LABA), long-acting muscarinic antagonists (LAMA) and inhaled corticosteroids are frequently prescribed to patients with COPD (1).

Some literature has associated anticholinergic use with increased risk of MCI (83) and dementia (83-85), and faster cognitive decline (86). However, participants in these studies used systemic rather than inhaled medication. Inhaled medication is much more targeted and will therefore have much less systemic effects. For example, tiotropium cannot cross the blood-brain barrier (87) which might suggest no or limited effect on cognitive functioning. However, no studies have investigated the effects of tiotropium on cognition yet.

The effects of corticosteroids or glucocorticoids (GC) on cognition have not been investigated in patients with COPD yet either. In general, mildly elevated GC levels improve cognition (88), but long-term administration can cause 'steroid dementia', characterized by impairments in episodic, declarative and working memory and executive function, and associated hippocampal and prefrontal dysfunction (89). Steroid dementia can appear within weeks of commencing GC treatment and is largely reversible upon its termination, although impairments can remain for years after termination (89, 90).

It is doubtful whether steroid dementia is a real risk in COPD. The recommended dose and duration for patients with an acute exacerbation (30-40 mg of prednisone daily for 7 to 14 days, where 5 days may be equally effective) (91) are much lower and shorter than those reported to cause steroid dementia (i.e., 60 mg/day for 7 months (90) or 40-60 mg/day for 37 days (92)).



In conclusion, to date, evidence that respiratory medication can contribute to impaired cognition in patients with COPD is lacking.

### *Exacerbations*

During acute exacerbations the above-mentioned determinants intensify and converge. Interestingly, exacerbations have an additional detrimental effect on cognition (22, 82, 92, 93). During acute exacerbations, levels of inflammation and cognitive functioning are inversely related (94), and as recovery from exacerbation-related CI seems very slow to non-existent, it can be speculated that regular COPD exacerbations can trigger a stepwise decrease in cognition. One study found no improvement in the 3 months after an exacerbation (22). However, because these patients were not cognitively tested prior to the exacerbation, it cannot be determined whether the exacerbation actually affected cognition in this study.

### *Comorbidities*

COPD often presents with comorbid conditions such as obstructive sleep apnea (OSA), depression and chronic heart failure (CHF), and a recent systematic review reported an increased risk prevalence of the metabolic syndrome in COPD compared to matched controls (95).

OSA contributes to decreased arterial oxygen saturation (96), causes sleep fragmentation and cortical and sympathetic arousal (96, 97), and affects attention, memory, psychomotor speed, visuospatial abilities, constructional abilities, executive functions and language abilities (96, 98). Hypoxemia, a shared component between OSA and advanced COPD, might underlie the cognitive deficits apparent in both diseases, namely attention, memory, executive function, psychomotor function and language abilities (98).

Depression is associated with decreased cognitive functioning (99) in COPD, but it probably only predicts 1-2% of the variation in cognition (3). A study investigating cognitive bias in patients with COPD and healthy controls with and without depression revealed that depressed patients with COPD showed a comparable pattern of bias compared to depressed healthy controls, whereas never-depressed patients showed much less bias (100). In conclusion, the influence of concurrent COPD and depression on cognition is still equivocal.

Many patients with COPD also suffer from CHF, and *vice versa* (17). The prevalence of CI in CHF is largely unknown, as prevalence estimates between 13.5% and 80% have been reported (17). CHF and COPD might have additive effects on cognition. Moreover, COPD and CHF are both associated with a high prevalence

of cerebrovascular diseases, which could lead to chronic cerebral hypoxia, impaired brain perfusion and ultimately brain damage and cognitive impairment (17). Furthermore, etiological similarities, such as cigarette smoking, may lead to a common set of symptoms, including CI (17).

Metabolic syndrome (MetS) is a cluster of metabolic risk factors, such as central obesity, dyslipidemia, hyperglycemia and dyslipidemia (95), with a prevalence of 34% in patients with COPD (95). MetS is strongly related to the risk of developing type 2 diabetes and cardiovascular disease (95) and has also been shown to have a deleterious influence on cognition (101, 102). However, some research also suggests that certain components of MetS have a larger effect than others (103, 104). It is yet unclear what the relative contribution of each of the components of MetS on cognition is.

### *Smoking*

Smoking is one of the largest risk factors for developing lung cancer or COPD, and it affects cognition in multiple ways. Firstly, it increases carbon monoxide and carbon dioxide levels in the blood, causing hypercapnia (105) and hypoxemia (106), respectively. Secondly, cigarette smoke contains many neurotoxic components, such as cadmium, nitric oxide and lead (107). Thirdly, the many free radicals in cigarette smoke are neurotoxic (108). And finally, chronic nicotine administration increases tau phosphorylation, a key component of Alzheimer's disease pathophysiology (107) and induced free radical production and depleted antioxidant levels in a rat model (109). Ultimately, all of these components cause decreased GM and WM volume and connectivity, and impair cognition (110-112).

Only one study investigated the influence of smoking on CI specifically in patients with COPD (113). Cognitive performance of patients with COPD was comparable to that of smokers, but both were significantly worse compared to normal reference values. The number of pack years and the duration of abstinence of the past smokers, consisting almost two-thirds of the sample, was not reported. Therefore, it cannot be determined with certainty to which degree smoking affects cognition beyond the effects of COPD. In general, smoking per se contributes to cognitive dysfunction, but there is also evidence of a relationship between impaired lung function and cognition independent of smoking (3, 114).

### *Dietary insufficiencies*

Diet and nutritional habits significantly impact on brain fitness, mental and cognitive health throughout life (115, 116). The relative abundance of specific dietary nutrients, depending on intake, bioavailability and metabolism, affects mental

health and cognitive ability via direct and indirect mechanisms that modulate neuronal function and synaptic plasticity (117). Chronic stress has been shown to negatively impact on brain plasticity and cognitive performance, for instance through the harmful effects of cortisol (118-120) and poor dietary habits are hypothesized to correlate with heightened stress reactivity and susceptibility (121) and greater cognitive decline in elderly (122). A healthy diet, rich in polyphenols, B vitamins, polyunsaturated fatty acids and dietary fibers exert favorable effects on cognitive performance, stress reactivity and neuroinflammation (117, 123). Unintended weight loss and muscle wasting is common in advanced COPD, but specific nutritional deficiencies that could affect cognition have received limited attention to date. However, next to disease severity, Collins *et al.* recently highlighted in a UK COPD population the importance of deprivation on malnutrition risk (124). An Australian study reported, next to low muscle mass, a high prevalence of deficiencies in vitamin D, vitamin B12 and iron in patients with COPD hospitalized with an acute exacerbation (125). A Dutch study investigating patients eligible for PR reported that vitamin D and calcium intake were below the recommended levels in more than 75% of patients, whereas vitamin A, C and E intakes were below the recommended levels in over one-third of patients (126). No studies have yet investigated the relationship between nutritional status or dietary pattern and cognitive performance in COPD.

### *Inactive lifestyle*

Higher levels of physical activity are associated with a reduced risk of cognitive decline and dementia (127). However, disease-related factors such as dyspnea and muscular metabolic abnormalities make it hard for many patients with COPD to be physically active. One systematic review found a mean daily step count of 2,237 (128), which is far less than the threshold for a low active (5,000 steps) or active lifestyle (10,000 steps) indicated by the same authors (128). Low physical activity levels and a sedentary lifestyle negatively impact on cognition in the general population (129, 130), and as such may contribute to CI in COPD as well.

Overall, disease-specific as well as lifestyle factors may contribute to the development of CI in COPD, but the relative potential synergistic contributions of the individual factors are yet unclear.

## Possible future interventions

### *Cognitive training*

Cognitive training can improve cognitive functioning in healthy elderly adults (131, 132), and can also improve some cognitive domains in those with MCI (131). Research into cognitive training specifically for patients with COPD is scarce. One trial attempted to ameliorate cognition in hypoxemic patients with COPD through an intervention aimed at improving attention, learning and logical-deductive thinking (133). During the first 2 weeks, the intervention group received group cognitive training, in the four weeks thereafter they received individual training, followed by home assignments two times per week. Booster sessions took place after 3 and 5 months. The control group received usual care and no cognitive training. Some cognitive domains improved in both the intervention and placebo group, but the intervention had no additional effect on cognition.

### *Exercise training*

Exercise can improve cognition through multiple pathways. It causes an increase in cerebral blood flow (38) and levels of cerebral growth factors such as brain-derived neurotrophic factor (134) and insulin growth factor-1 (135). These growth factors are involved in many functions which are important for cognition. They influence the rate of differentiation and apoptosis of cerebral cells (136), and regulate long-term potentiation (137) and hippocampal neurogenesis (136). Multiple previous studies have shown the benefit of exercise on cognition in COPD patients (138, 139), and in the study by Park *et al.* the 6-minute walking distance was the only potentially modifiable variable that was related to worsening cognitive functioning over time (27). The question remains, however, whether these improvements consolidate into the longer term if exercise is discontinued (11).

### *Smoking cessation*

Given the deleterious influences of smoking on cognition, it is unsurprising that smoking cessation is beneficial. However, its effects on cognitive functioning have only been assessed in the general population. It appears that the number of pack years negatively affects cognitive functioning, but also that cognitive functioning improves with longer duration of abstinence (140). This implies that abstinence pays off at any age, with the largest benefits coming to those who stop the earliest. Future research should further investigate the effects of smoking cessation on cognition in COPD in more detail.

### *Dietary intervention*

Dietary intervention may imply adopting a different dietary pattern or supplementing the habitual diet with specific nutrients. The Mediterranean diet, the Dietary Approach to Stop Hypertension (DASH) diet and the Mediterranean-DASH Intervention for Neurodegenerative Delay (MIND) diet have been proposed as beneficial to cognition (141-143). Drastic dietary changes, however, are not likely to be feasible for patients with advanced COPD.

These interventions imply that CI in COPD is not something that patients have to passively undergo but can be readily applied in daily life. However, more research is needed to assess the feasibility and efficacy of the individual interventions or multimodalities thereof.

### **Conclusion**

CI in patients with COPD is a problem with a high prevalence and large consequences, yet it is still under-recognized and under-investigated. More research aimed at unraveling the etiology and appropriate interventions to diminish cognitive decline or treat CI in patients with COPD is needed to benefit the patients as well as their loved ones.

Caregivers should pay more attention to potential CI in their patients, as CI may have large consequences on self-management and health outcomes. In a clinical context, administering a brief screening tool may help in identifying patients who need referral to a specialist for further investigation. These patients also need more time and attention, for instance while making sure they understand how to properly use their medication.

### **Expert commentary**

Technical developments in the field of cognitive neuroscience enable more detailed insight in the *pathophysiology* of CI in COPD. Increasing magnetic resonance imaging resolution allows a more detailed picture of the brain, techniques such as diffusion tensor imaging allow focusing specifically on white matter instead of the brain as a whole, and shorter and more effective scanning protocols can make brain imaging more accessible and affordable.

Investigating novel research directions might also be worthwhile. For instance, the synthesis and proper functioning of many neurotransmitters, including the catecholamines, glutamate, aspartate and perhaps most importantly acetylcholine (as it is widely available in the brain but also the most important neurotransmitter in the airways) depend on oxygen availability (144). Hypoxia-based neurotransmitter abnormalities might therefore constitute a third type of cerebral abnormality contributing to CI, next to structural and functional abnormalities. However, this has received scarce research attention to date in the general population, and especially in COPD.

Longitudinal studies are essential for understanding the *causes and consequences* of CI in COPD. They can elucidate the development of cognitive decline in COPD in relation to the development of other potentially relevant variables such as impaired lung function, hypoxia or inflammation. In this way, the relative contributions of disease- and aging-related factors can also be further disentangled.

Interesting *treatment options* for CI in COPD include the potential role of specific nutrients in ameliorating cognition. The Mediterranean diet, which is characterized by a high intake of plant-based foods, moderate-to-high fish and seafood consumption and scarce use of dairy products and meat (141), has earlier been shown to improve cognition in a randomized controlled trial (141). Furthermore, polyunsaturated fatty acids and polyphenols can ameliorate cognition and have a positive effect on various neurobiological processes (145). Also, testing potential synergistic effects of combinations of interventions could be worthwhile, such as exercise training and cognitive training.

An important prerequisite for all the above research questions is the choice of the right cognitive test instruments. Often applied screening tools such as the MMSE or MoCA can be useful in a clinical setting to identify patients in need of further neuropsychological assessment and therapy, but cannot give a detailed overview of the exact nature of a person's deficits. Larger, well-defined and evidence-based test batteries are essential in order to get a comprehensive overview of a person's neuropsychological functioning (146).

## Five-year view

In conclusion, it is important to further investigate the pathophysiology, causes and consequences of CI in COPD in the next five years, and to develop tailored intervention strategies. Developments in the fields of cognitive neuroscience and neuropsychology will enable a more detailed picture of the pathophysiology of

CI in COPD. Longitudinal studies can pinpoint its determinants and identify its consequences, and much work remains to be done in finding (the most) effective treatments.

## References

1. Global Initiative for Chronic Obstructive Lung Disease (GOLD). Global Strategy for the Diagnosis, Management and Prevention of Chronic Obstructive Pulmonary Disease (2018 Report). Agusti A, Vogelmeier C, editors. 2018.
2. Cavaillès A, Brinchault-Rabin G, Dixmier A, Goupil F, Gut-Gobert C, Marchand-Adam S, *et al.* Comorbidities of COPD. *European Respiratory Review*. 2013;22:454-75.
3. Dodd JW, Getov SV, Jones PW. Cognitive function in COPD. *European Respiratory Journal*. 2010;35:913-22.
4. Cleutjens FA, Janssen DJ, Ponds RW, Dijkstra JB, Wouters EF. COgnitive-Pulmonary Disease. *BioMed Research International*. 2014:697825.
5. Lezak MD, Howieson DB, Bigler ED, Tranel D. *Neuropsychological Assessment* (5th edition). New York, NY: Oxford University Press; 2012.
6. Franz EA, Gillett G. John Hughlings Jackson's evolutionary neurology: a unifying framework for cognitive neuroscience. *Brain*. 2011;134(10):3114-20.
7. Diamond A. Executive functions. *Annual Review of Psychology*. 2013;64:135-68.
8. Kolb B, Whishaw IQ. *An introduction to brain and behavior* (3rd edition). New York, NY: Worth Publishers; 2011.
9. Petersen RC. Mild cognitive impairment. *Continuum*. 2016;22(2):404-18.
10. Belleville S, Fouquet C, Hudon C, Vignon Zomahoun HT, Croteau J. Neuropsychological measures that predict progression from mild cognitive impairment to Alzheimer's type dementia in older adults: a systematic review and meta-analysis. *Neuropsychological Review*. 2017;27(4):328-53.
11. Ouellette DR, Lavoie KL. Recognition, diagnosis and treatment of cognitive and psychiatric disorders in patients with COPD. *International Journal of COPD*. 2017;12:639-50.
12. Zhang H, Wang X, Lin J, Sun Y, Huang Y, Yang T, *et al.* Grey and white matter abnormalities in chronic obstructive pulmonary disease: a case-control study. *BMJ Open*. 2012;2:e000844.
13. Schou L, Ostergaard B, Rasmussen LS, Rydahl-Hansen S, Phanareth K. Cognitive dysfunction in patients with chronic obstructive pulmonary disease - a systematic review. *Respiratory Medicine*. 2012;106:1071-81.
14. Zhang J, Chen J, Yu Q, Fan C, Zhang R, Lin J, *et al.* Alteration of spontaneous brain activity in COPD patients. *International Journal of COPD*. 2016;11:1713-9.
15. Zhang H, Wang X, Lin J, Sun Y, Huang Y, Yang T, *et al.* Reduced regional gray matter volume in patients with chronic obstructive pulmonary disease: a voxel-based morphometry study. *American Journal of Neuroradiology*. 2013;34(2):334-9.
16. Andrianopoulos V, Gloeckl R, Vogiatzis I, Kenn K. Cognitive impairment in COPD: should cognitive evaluation be part of respiratory assessment? *Breathe*. 2017;13(1):e1-e9.



17. Yohannes AM, Chen W, Moga AM, Leroi I, Connolly MJ. Cognitive impairment in chronic obstructive pulmonary disease and chronic heart failure: a systematic review and meta-analysis of observational studies. *JAMDA*. 2017;18:451.e1-.e11.
18. Cleutjens FAHM, Franssen F, Spruit MA, Vanfleteren LEGW, Gijssen C, Dijkstra JB, et al. Domain-specific cognitive impairment in patients with COPD and control subjects. *International Journal of Chronic Obstructive Pulmonary Disease*. 2017;12:1-11.
19. Cleutjens FAHM, Spruit MA, Ponds RWHM, Vanfleteren LEGW, Franssen F, Gijssen C, et al. Cognitive impairment and clinical characteristics in patients with chronic obstructive pulmonary disease. *Chronic Respiratory Disease*. 2018;15(2):91-102.
20. Pierobon A, Bottelli ES, Ranzini L, Bruschi C, Maestri R, Bertolotti G, et al. COPD patients' self-reported adherence, psychosocial factors and mild cognitive impairment in pulmonary rehabilitation. *International Journal of COPD*. 2017;12:2059-67.
21. Roncero C, Campuzano AI, Quintano JA, Molina J, Perez J, Miravittles M. Cognitive status among patients with chronic obstructive pulmonary disease. *International Journal of COPD*. 2016;11:543-51.
22. Dodd JW, Charlton RA, Van den Broek MD, Jones PW. Cognitive dysfunction in patients hospitalized with acute exacerbation of COPD. *Chest*. 2013;144(1):119-27.
23. Fekri MS, Hashemi-Bajgani S-M, Naghibzadeh-Tahami A, Arabnejad F. Cognitive impairment among patients with chronic obstructive pulmonary disease compared to normal individuals. *Tanaffos*. 2017;16(1):34-9.
24. Jolles J, Van Boxtel MPJ, Ponds RW, Metsemakers JF, Houx PJ. The Maastricht Aging Study (MAAS) - the longitudinal perspective of cognitive aging. *Tijdschrift voor Gerontologie en Geriatrie*. 1998;29(3):120-9.
25. López-Torres I, Valenza MC, Torres-Sánchez IC-M, I., Rodríguez-Torres J, Moreno-Ramírez MP. Changes in cognitive status in COPD patients across clinical stages. *COPD*. 2016;13(3):327-32.
26. Rusanen M, Ngandu T, Laatikainen T, Tuomilehto J, Soininen H, Kivipelto M. Chronic obstructive pulmonary disease and asthma and the risk of mild cognitive impairment and dementia: a population based CAIDE study. *Current Alzheimer Research*. 2013;10:549-55.
27. Park SK. Trajectories of change in cognitive function in people with COPD. *Journal of Clinical Nursing*. 2018;27(7-8):1529-42.
28. Korfiatis S, Stranjalis GP, A., Psachoulia C, Daskalakis G, Antsaklis A, Sakas DE. Serum S-100B protein as a biochemical marker of brain injury: a review of current concepts. *Current Medicinal Chemistry*. 2006;13(30):3719-31.
29. Esser RW, Stoeckel MC, Kirsten A, Watz H, Taube K, Lehmann K, et al. Structural brain changes in patients with COPD. *Chest*. 2016;149(2):426-34.

30. Spilling CA, Jones PW, Dodd JW, Barrick TR. White matter lesions characterise brain involvement in moderate to severe chronic obstructive pulmonary disease, but cerebral atrophy does not. *BMC Pulmonary Medicine*. 2017;17:92.
31. Dodd JW, Chung AW, Van den Broek MD, Barrick TR, Charlton RA, Jones PW. Brain structure and function in chronic obstructive pulmonary disease - a multimodal cranial magnetic resonance imaging study. *American Journal of Respiratory and Critical Care Medicine*. 2012;186(3):240-5.
32. Ryu C-W, Jahng G-H, Choi CW, Rhee HY, Kim M-J, Kim S-M, *et al*. Microstructural change of the brain in chronic obstructive pulmonary disease: a voxel-based investigation by MRI. *Journal of Chronic Obstructive Pulmonary Disease*. 2013;10:357-66.
33. Cleutjens FA, Ponds RW, Spruit MA, Burgmans S, Jacobs HIL, Gronenschild EHBM, *et al*. The relationship between cerebral small vessel disease, hippocampal volume and cognitive functioning in patients with COPD: an MRI study. *Frontiers in Aging Neuroscience*. 2017;9:88.
34. Chen J, Lin I-T, Zhang H, Lin J, Zheng S, Fan M, *et al*. Reduced cortical thickness, surface area in patients with chronic obstructive pulmonary disease: a surface-based morphometry and neuropsychological study. *Brain Imaging and Behavior*. 2016;10:464-76.
35. Barbey AK, Koenigs M, Grafman J. Dorsolateral prefrontal contributions to human working memory. *Cortex*. 2013;49:1195-205.
36. Wang C, Ding Y, Shen B, Gao D, An J, Peng K, *et al*. Altered gray matter volume in stable chronic obstructive pulmonary disease with subclinical cognitive impairment: an exploratory study. *Neurotoxicity Research*. 2017;31(4):453-63.
37. Lövlblad KO, Schaller K, Vargas MI. The fornix and limbic system. *Seminars in Ultrasound, CT, and MR*. 2014;35(5):459-73.
38. Sachdev PS, Anstey KJ, Parslow RA, Wen W, Maller J, Kumar R, *et al*. Pulmonary function, cognitive impairment and brain atrophy in a middle-aged community sample. *Dementia and Geriatric Cognitive Disorders*. 2006;21:300-8.
39. Taki Y, Kinomura S, Ebihara S, Thyreau B, Sato K, Goto R, *et al*. Correlation between pulmonary function and brain volume in healthy elderly subjects. *Neuroradiology*. 2013;55(6):689-95.
40. Poulouse SM, Miller MG, Scott T, Shukitt-Hale B. Nutritional factors affecting adult neurogenesis and cognitive function. *Advances in Nutrition*. 2017;8:804-11.
41. Li J, Fei GH. The unique alterations of hippocampus and cognitive impairment in chronic obstructive pulmonary disease. *Respiratory Research*. 2013;14:140.
42. Borson S, Scanlan J, Friedman S, Zuhr E, Fields J, Aylward E, *et al*. Modeling the impact of COPD on the brain. *International Journal of COPD*. 2008;3(3):429-34.

43. Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME. The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *PNAS*. 2005;102(27):9673-8.
44. Hu X, Wang H, Tu Y, Fei M, Yin M, Fei G, *et al*. Alterations of the default mode network and cognitive impairments in patients with chronic obstructive pulmonary disease. *International Journal of COPD*. 2018;13:519-28.
45. Leung JM, Sin DD. Chronic obstructive pulmonary disease and stroke: strange bedfellows. *American Journal of Respiratory and Critical Care Medicine*. 2016;193(3):227-8.
46. Portegies MLP, Lahousse L, Joos GF, Hofman A, Koudstaal PJ, Stricker BH, *et al*. Chronic obstructive pulmonary disease and the risk of stroke: the Rotterdam Study. *American Journal of Respiratory and Critical Care Medicine*. 2016;193(3):251-8.
47. Soderholm M, Inghammar M, Hedblad B, Egesten A, Engstrom G. Incidence of stroke and stroke subtypes in chronic obstructive pulmonary disease. *European Journal of Epidemiology*. 2016;31(2):159-68.
48. Morgan AD, Sharma C, Rothnie KJ, Potts J, Smeeth L, Quint JK. Chronic obstructive pulmonary disease and the risk of stroke. *Annals of the American Thoracic Society*. 2017;14(5):754-65.
49. Kim YR, Hwang IC, Lee YJ, Ham EB, Park DK, Kim S. Stroke risk among patients with chronic obstructive pulmonary disease: a systematic review and meta-analysis. *Clinics*. 2018;73:e177.
50. Effing TW, Vercoulen JH, Bourbeau J, Trappenburg J, Lenferink A, Cafarella P, *et al*. Definition of a COPD self-management intervention: International Expert Group consensus. *European Respiratory Journal*. 2016;48(1):46-54.
51. Baird C, Lovell J, Johnson M, Shiell K, Ibrahim JE. The impact of cognitive impairment on self-management in chronic obstructive pulmonary disease: a systematic review. *Respiratory Medicine*. 2017;129:130-9.
52. Dulohery MM, Schroeder DR, Benzo R. Cognitive function and living situation in COPD: is there a relationship with self-management and quality of life? *International Journal of COPD*. 2015;10:1883-9.
53. Benzo R, Kirsch JL, Dulohery MM, Abascal-Bolado B. Emotional intelligence: a novel outcome associated with wellbeing and self-management in chronic obstructive pulmonary disease. *Annals of the American Thoracic Society*. 2016;13(1):10-6.
54. Meek P, Lareau SC, Anderson D. Memory for symptoms in COPD patients: how accurate are their reports? *European Respiratory Journal*. 2001;18(3):474-81.
55. Allen SC, Ragab S. Ability to learn inhaler technique in relation to cognitive scores and tests of praxis in old age. *Postgraduate Medical Journal*. 2002;78:37-9.

56. Brega AG, Grigsby J, Kooken R, Hamman RF, Baxter J. The impact of executive cognitive functioning on rates of smoking cessation in the San Luis Valley Health and Aging Study. *Age and Ageing*. 2008;37(5):521-5.
57. Fox AT, Martin LE, Bruce J, Moreno JL, Staggs VS, Lee HS, *et al*. Executive function fails to predict smoking outcomes in a clinical trial to motivate smokers to quit. *Drug and Alcohol Dependence*. 2017;175:227-31.
58. Cohen-Mansfield J. Predictors of smoking cessation in old-old age. *Nicotine and Tobacco Research*. 2016;18(7):1675-9.
59. Ozyemisci-Taskiran O, Bozkurt SO, Kokturk N, Karatas GK. Is there any association between cognitive status and functional capacity during exacerbation of chronic obstructive pulmonary disease? *Chronic Respiratory Disease*. 2015;12(3):247-55.
60. Yazar EE, Aydin S, Gunluoglu G, Kamat S, Gungen AC, Yildiz P. Clinical effects of cognitive impairment in patients with chronic obstructive pulmonary disease. *Chronic Respiratory Disease*. 2018;15(3):306-14.
61. Cleutjens FAHM, Spruit MA, Ponds RWHM, Vanfleteren LEGW, Franssen FM, Dijkstra JB, *et al*. The impact of cognitive impairment on efficacy of pulmonary rehabilitation in patients with COPD. *JAMDA*. 2017;18(5):420-6.
62. Schure MB, Borson S, Nguyen HQ, Trittschuh EH, Thielke SM, Pike KC, *et al*. Associations of cognition with physical functioning and health-related quality of life among COPD patients. *Respiratory Medicine*. 2016;114:46-52.
63. Martinez CH, Richardson CR, Han MK, Cigolle CT. Chronic obstructive pulmonary disease, cognitive impairment, and development of disability: the health and retirement study. *Annals of the American Thoracic Society*. 2014;11(9):1362-70.
64. Chang SS, Chen S, McAvay GJ, Tinetti ME. Effect of coexisting chronic obstructive pulmonary disease and cognitive impairment on health outcomes in older adults. *Journal of the American Geriatric Society*. 2012;60(10):1839-46.
65. Dodd JW, Novotny P, Sciruba FC, Benzo R, NETT Research Group. Executive function, survival, and hospitalization in chronic obstructive pulmonary disease: a longitudinal analysis of the National Emphysema Treatment Trial (NETT). *Annals of the American Thoracic Society*. 2015;12(10):1473-81.
66. Antonelli Incalzi R, Corsonello A, Pedone C, Trojano L, Acanfora D, Spada A, *et al*. Drawing impairment predicts mortality in severe COPD. *Chest*. 2006;130(6):1687-94.
67. Yohannes AM, Raue PJ, Kanellopoulos D, McGovern A, Sirey JA, Kiosses DN, *et al*. Predictors of all-cause mortality in patients with severe COPD and major depression admitted to a rehabilitation hospital. *Chest*. 2016;149(2):467-73.
68. Alexandre F, Heraud N, Varray A. Is nocturnal desaturation a trigger for neuronal damage in chronic obstructive pulmonary disease? *Medical Hypotheses*. 2015;84:25-30.

69. Barnes PJ. Chronic obstructive pulmonary disease: effects beyond the lungs. *PLoS Medicine*. 2010;7(3):e1000220.
70. Antonelli Incalzi R, Marra C, Giordano A, Calcagni ML, Cappa A, Basso S, *et al*. Cognitive impairment in chronic obstructive pulmonary disease - a neuropsychological and SPECT study. *Journal of Neurology*. 2003;250(3):325-32.
71. Ortapamuk H, Naldoken S. Brain perfusion abnormalities in chronic obstructive pulmonary disease: comparison with cognitive impairment. *Annals of Nuclear Medicine*. 2006;20(2):99-106.
72. Teppema LJ, Dahan A. The ventilatory response to hypoxia in mammals: mechanisms, measurement, and analysis. *Physiology Reviews*. 2010;90(2):675-754.
73. Sicard KM, Duong TQ. Effects of hypoxia, hyperoxia, and hypercapnia on baseline and stimulus-evoked BOLD, CBF, and CMRO<sub>2</sub> in spontaneously breathing animals. *NeuroImage*. 2005;25(3):850-8.
74. Sumiyoshi A, Suzuki H, Shimokawa H, Kawashima R. Neurovascular uncoupling under mild hypoxic hypoxia: an EEG-fMRI study in rats. *Journal of Cerebral Blood Flow and Metabolism*. 2012;32(10):1853-8.
75. Marshall O, Uh J, Lurie D, Lu H, Milham MP, Ge Y. The influence of mild carbon dioxide on brain functional homotopy using resting-state fMRI. *Human Brain Mapping*. 2015;36(10):3912-21.
76. Hartmann SE, Pialoux V, Leigh R, Poulin MJ. Decreased cerebrovascular response to CO<sub>2</sub> in post-menopausal females with COPD: role of oxidative stress. *European Respiratory Journal*. 2012;40:1354-61.
77. Nguyen JC, Killcross AS, Jenkins TA. Obesity and cognitive decline: role of inflammation and vascular changes. *Frontiers in Neuroscience*. 2014;8:375.
78. Solas M, Milagro FI, Ramirez MJ, Martinez JA. Inflammation and gut-brain axis link obesity to cognitive dysfunction: plausible pharmacological interventions. *Current Opinion in Pharmacology*. 2017;37:87-92.
79. Siervo M, Harrison SL, Jagger C, Robinson L, Stephan BCM. Metabolic syndrome and longitudinal changes in cognitive function: a systematic review and meta-analysis. *Journal of Alzheimer's Disease*. 2014;41:151-61.
80. Panza F, Frisardi VC, C., Imbimbo BP, Vendemiale G, Santamato A, D'Onofrio G, *et al*. Metabolic syndrome and cognitive impairment: current epidemiology and possible underlying mechanisms. *Journal of Alzheimer's disease*. 2010;21(3):691-724.
81. Schram MT, Euser SM, De Craen AJ, Witteman JC, Frölich M, Hofman A, *et al*. Systemic markers of inflammation and cognitive decline in old age. *Journal of the American Geriatric Society*. 2007;55(5):708-16.
82. Crisan AF, Oancea C, Timar B, Fira-Mladinescu O, Crisan A, Tudorache V. Cognitive impairment in chronic obstructive pulmonary disease. *PLoS One*. 2014;9(7):e102468.

83. Pfistermeister B, Tümena T, Gaßmann K-G, Maas R, Fromm MF. Anticholinergic burden and cognitive function in a large German cohort of hospitalized geriatric patients. *PLoS One*. 2017;12(2):e0171353.
84. Jessen F, Kaduszkiewicz H, Daerr M, Bickel H, Pentzek M, Riedel-Heller S, et al. Anticholinergic drug use and risk for dementia: target for dementia prevention. *European Archives of Psychiatry and Clinical Neuroscience*. 2010;260(Suppl 2):S111-S5.
85. Gray SL, Anderson ML, Dublin S, Hanlon JT, Hubbard R, Walker R, et al. Cumulative use of strong anticholinergics and incident dementia: a prospective cohort study. *JAMA Internal Medicine*. 2015;175(3):401-7.
86. Fox C, Richardson K, Maidment ID, Savva GM, Matthews FE, Smithard D, et al. Anticholinergic medication use and cognitive impairment in the older population: the Medical Research Council cognitive function and ageing study. *Journal of the American Geriatric Society*. 2011;59:1477-83.
87. Heredia JL. Tiotropium bromide: an update. *Open Respiratory Medicine Journal*. 2009;3:43-52.
88. Joels M. Corticosteroid effects in the brain: U-shape it. *Trends in Pharmacological Sciences*. 2006;27(5):244-50.
89. Wolkowitz OM, Lupien SJ, Bigler ED. The "steroid dementia syndrome": a possible model of human glucocorticoid neurotoxicity. *Neurocase*. 2007;13(3):189-200.
90. Cipriani G, Picchi L, Vedovello M, Nuti A. Reversible dementia from corticosteroid therapy. *Clinical Geriatrics*. 2012;20(7):3841.
91. Walters JAE, Tan DJ, White CJ, Wood-Baker R. Different durations of corticosteroid therapy for exacerbations of chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews*. 2018;3(3):CD006897.
92. Zhang X, Cai X, Shi X, Zheng Z, Zhang A, Guo J, et al. Chronic obstructive pulmonary disease as a risk factor for cognitive dysfunction: a meta-analysis of current studies. *Journal of Alzheimer's Disease*. 2016;52(1):101-11.
93. Tulek B, Atalay NB, Yildirim G, Kanat F, Suerdem M. Cognitive function in chronic obstructive pulmonary disease: relationship to global initiative for chronic obstructive lung disease 2011 categories. *Respirology*. 2014;19(6):873-80.
94. Wedzicha JA, Seemungal TAR. COPD exacerbations: defining their cause and prevention. *Lancet*. 2007;370(9589):786-96.
95. Cebron Lipovec N, Beijers RJ, Van den Borst B, Doehner W, Lainscak M, Schols AMWJ. The prevalence of metabolic syndrome in chronic obstructive pulmonary disease: a systematic review. *COPD*. 2016;13(3):399-406.
96. Andreou G, Vlachos F, Mekanikas K. Effects of chronic obstructive pulmonary disease and obstructive sleep apnea on cognitive functions: evidence for a common nature. *Sleep Disorders*. 2014;2014:768210.

97. Daulatzai MA. Pathogenesis of cognitive dysfunction in patients with obstructive sleep apnea: a hypothesis with emphasis on the nucleus tractus solitarius. *Sleep Disorders*. 2012;2012:251096.
98. Olaithe M, Bucks RS, Hillman DR, Eastwood PR. Cognitive deficits in obstructive sleep apnea: insights from a meta-review and comparison with deficits observed in COPD, insomnia, and sleep deprivation. *Sleep Medicine Reviews*. 2018;38:39-49.
99. Aras YG, Tunc A, Gungen BD, Gungen AC, Aydemir Y, Demiyurek BE. The effects of depression, anxiety and sleep disturbances on cognitive impairment in patients with chronic obstructive pulmonary disease. *Cognitive Neurodynamics*. 2017;11(6):565-71.
100. Fritzsche A, Watz H, Magnussen H, Tuinmann G, Lowe B, Von Leupoldt A. Cognitive biases in patients with chronic obstructive pulmonary disease and depression - a pilot study. *British Journal of Health Psychology*. 2013;18:827-43.
101. Viscogliosi G, Donfrancesco C, Palmieri L, Giampaoli S. The metabolic syndrome and 10-year cognitive and functional decline in very old men - a population-based study. *Archives of Gerontology and Geriatrics*. 2017;70:62-6.
102. Alfaro FJ, Lioutas VA, Pimentel DA, Chung CC, Bedoya F, Yoo WK, *et al*. Cognitive decline in metabolic syndrome is linked to microstructural white matter abnormalities. *Journal of Neurology*. 2016;263(12):2505-14.
103. Assuncao N, Sudo FK, Drummond C, De Felice FG, Mattos P. Metabolic syndrome and cognitive decline in the elderly: a systematic review. *PLoS One*. 2018;13(3):e0194990.
104. Overman MJ, Pendleton N, O'Neill TW, Bartfai G, Casanueva FF, Forti G, *et al*. Glycemia but not the metabolic syndrome is associated with cognitive decline: findings from the European Male Ageing Study. *American Journal of Geriatric Psychiatry*. 2017;25(6):662-71.
105. Grant I, Heaton RK, McSweeney AJ, Adams KM, Timms RM. Neuropsychologic findings in hypoxemic chronic obstructive pulmonary disease. *Archives of Internal Medicine*. 1982;142(8):1470-6.
106. Ling J, Heffernan T. The cognitive deficits associated with second-hand smoking. *Frontiers in Psychiatry*. 2016;7:46.
107. Swan GE, Lessov-Schlaggar CN. The effects of tobacco smoke and nicotine on cognition and the brain. *Neuropsychological Review*. 2007;17:259-73.
108. Waisman Campos M, Serebrisky D, Castaldelli-Maia JM. Smoking and cognition. *Current Drug Abuse Reviews*. 2016;9:1-4.
109. Qiao D, Seidler FJ, Slotkin TA. Oxidative mechanisms contributing to the developmental neurotoxicity of nicotine and chlorpyrifos. *Toxicology and Applied Pharmacology*. 2005;206(1):17-26.

110. Zhang X, Salmeron BJ, Ross TJ, Gu H, Geng X, Yang Y, *et al.* Anatomical differences and network characteristics underlying smoking cue reactivity. *NeuroImage*. 2011;54(1):131-41.
111. Zhang X, Salmeron BJ, Ross TJ, Geng X, Yang Y, Stein EA. Factors underlying prefrontal and insula structural alterations in smokers. *NeuroImage*. 2011;54(1):42-8.
112. Yu R, Zhao L, Lu L. Regional grey and white matter changes in heavy male smokers. *PLoS One*. 2011;6(11):e27440.
113. Dal Negro RW, Bonadiman L, Tognella S, Bricolo FP, Turco P. Extent and prevalence of cognitive dysfunction in chronic obstructive pulmonary disease, chronic non-obstructive bronchitis, and in asymptomatic smokers, compared to normal reference values. *International Journal of COPD*. 2014;26(9):675-83.
114. Torres-Sanchez I, Rodriguez-Alzueta E, Cabrera-Martos I, Lopez-Torres I, Moreno-Ramírez MP, Valenza MC. Cognitive impairment in COPD: a systematic review. *Jornal Brasileiro de Pneumologia*. 2015;41(2):182-90.
115. Lai JS, Hiles S, Bisquera A, Hure AJ, McEvoy M, Attia J. A systematic review and meta-analysis of dietary patterns and depression in community-dwelling adults. *American Journal of Clinical Nutrition*. 2014;99(1):181-97.
116. Abbatecola AM, Russo M, Barbieri M. Dietary patterns and cognition in older persons. *Current Opinion in Clinical Nutrition and Metabolic Care*. 2018;21(1):10-3.
117. Bazinet RP, Layé S. Polyunsaturated fatty acids and their metabolites in brain function and disease. *Nature Reviews Neuroscience*. 2014;15(12):771-85.
118. Radley J, Morilak D, Viau V, Campeau S. Chronic stress and brain plasticity: mechanisms underlying adaptive and maladaptive changes and implications for stress-related CNS disorders. *Neuroscience and Biobehavioral Reviews*. 2015;58:79-91.
119. Marin M-F, Lord C, Andrews J, Juster R-P, Sindi Sh, Arsenault-Lapierre G, *et al.* Chronic stress, cognitive functioning and mental health. *Neurobiology of Learning and Memory*. 2011;96(4):583-95.
120. Sandi C. Stress and cognition. *Wiley Interdisciplinary Reviews - Cognitive Sciences*. 2013;4(3):245-61.
121. Allen AP, Dinan TG, Clarke G, Cryan JF. A psychology of the human brain-gut-microbiome axis. *Social and Personality Psychology Compass*. 2017;11(4):e12309.
122. Diet and nutrition in dementia and cognitive decline. London, England: Academic Press; 2015.
123. Smyth A, Dehghan M, O'Donnell M, Anderson C, Teo K, Gao P, *et al.* Healthy eating and reduced risk of cognitive decline: a cohort from 40 countries. *Neurology*. 2015;84(22):2258-65.
124. Collins PF, Elia M, Kurukulaaratchy RJ, Stratton RJ. The influence of deprivation on malnutrition risk in outpatients with chronic obstructive pulmonary disease (COPD). *Clinical Nutrition*. 2018;37(1):144-8.



125. Horadagoda C, Dinihan T, Roberts M, Kairaitis K. Body composition and micronutrient deficiencies in patients with an acute exacerbation of chronic obstructive pulmonary disease. *Internal Medicine Journal*. 2017;47(9):1057-63.
126. Van de Bool C, Mattijssen-Verdonschot C, Van Melick PPMJ, Spruit MA, Franssen FME, Wouters EFM, *et al*. Quality of dietary intake in relation to body composition in patients with chronic obstructive pulmonary disease eligible for pulmonary rehabilitation. *European Journal of Clinical Nutrition*. 2014;68(2):159-65.
127. Blondell SJ, Hammersley-Mather R, Veerman JL. Does physical activity prevent cognitive decline and dementia? A systematic review and meta-analysis of longitudinal studies. *BMC Public Health*. 2014;14:510.
128. Tudor-Locke C, Craig CL, Aoyagi Y, Bell RC, Croteau KA, De Bourdeaudhuij I, *et al*. How many steps/day are enough? For older adults and special populations. *International Journal of Behavioral Nutrition and Physical Activity*. 2011;8:80.
129. Kobayashi Y, Takahashi Y, Seki T, Kaneta T, Amarume K, Kasai M, *et al*. Decreased physical activity associated with executive dysfunction correlates with cognitive impairment among older adults in the community: a retrospective analysis from the Kurihara project. *Dementia and Geriatric Cognitive Disorders Extra*. 2016;6(2):350-60.
130. Falck RS, Davis JC, Liu-Ambrose T. What is the association between sedentary behaviour and cognitive function? A systematic review. *British Journal of Sports Medicine*. 2017;51(10):800-11.
131. Cinzia G, Roberta P, Fabrizia L, Demetrio P. The effects of cognitive training for elderly: results from My Mind project. *Rejuvenation Research*. 2016;19(6):485-94.
132. Shao YK, Mang J, Li PL, Wang J, Deng T, Xu ZX. Computer-based cognitive programs for improvement of memory, processing speed and executive function during age-related cognitive decline: a meta-analysis. *PLoS One*. 2015;10(6):e0130831.
133. Antonelli Incalzi R, Corsonello A, Trojano L, Pedone C, Acanfora D, Spada A, *et al*. Cognitive training is ineffective in hypoxemic COPD: a six-month randomized controlled trial. *Rejuvenation Research*. 2008;11:239-50.
134. Szuhany KL, Bugatti M, Otto MW. A meta-analytic review of the effects of exercise on brain-derived neurotrophic factor. *Journal of Psychiatric Research*. 2015;60:56-64.
135. Chen HT, Chung YC, Chen YJ, Ho SY, Wu HJ. Effects of different types of exercise on body composition, muscle strength, and IGF-1 in the elderly with sarcopenic obesity. *Journal of the American Geriatric Society*. 2017;65(4):827-32.
136. Pasco JA, Williams LJ, Jacka FN, Stupka N, Brennan-Olsen SL, Holloway KL, *et al*. Sarcopenia and the common mental disorders: a potential regulatory role of skeletal muscle on brain function? *Current Osteoporosis Reports*. 2015;13:351-7.
137. Leal G, Afonso PM, Salazar IL, Duarte CB. Regulation of hippocampal synaptic plasticity by BDNF. *Brain Research*. 2015;1621:82-101.

138. Aquino G, Iuliano E, Di Cagno A, Vardaro A, Fiorilli G, Moffa S, *et al.* Effects of combined training vs aerobic training on cognitive functions in COPD: a randomized controlled trial. *International Journal of COPD*. 2016;11:711-8.
139. Etnier JL, Berry M. Fluid intelligence in an older COPD sample after short- or long-term exercise. *Medicine and Science in Sports and Exercise*. 2001;33(10):1620-8.
140. Mons U, Schottker B, Muller H, Kliegel M, Brenner H. History of lifetime smoking, smoking cessation and cognitive function in the elderly population. *European Journal of Epidemiology*. 2013;28(10):823-31.
141. Martínez-Lapiscina EH, Clavero P, Toledo E, Estruch R, Salas-Salvadó J, San Julián B, *et al.* Mediterranean diet improves cognition: the PREDIMED-NAVARRA randomised trial. *Journal of Neurology, Neurosurgery and Psychiatry*. 2013;84(12):1318-25.
142. McEvoy CT, Guyer H, Langa KM, Yaffe K. Neuroprotective diets are associated with better cognitive function: the Health and Retirement Study. *Journal of the American Geriatric Society*. 2017;65(8):1857-62.
143. Tangney CC, Li H, Wang Y, Barnes L, Schneider JA, Bennett DA, *et al.* Relation of DASH- and Mediterranean-like dietary patterns to cognitive decline in older persons. *Neurology*. 2014;83(16):1410-6.
144. Gibson GE, Pulsinelli W, Blass JP, Duffy TE. Brain dysfunction in mild to moderate hypoxia. *American Journal of Medicine*. 1981;70(6):1247-54.
145. Spencer SJ, Korosi A, Layé S, Shukitt-Hale B, Barrientos RM. Food for thought: how nutrition impacts cognition and emotion. *NPJ Science of Food*. 2017;1:7.
146. Roebuck-Spencer TM, Glen T, Puente AE, Denney RL, Ruff RM, Hostetter G, *et al.* Cognitive screening tests versus comprehensive neuropsychological test batteries: a National Academy of Neuropsychology education paper. *Archives of Clinical Neuropsychology*. 2017;32(4):491-8.



# Chapter 5

## Cognitive performance in relation to metabolic disturbances in patients with COPD

Martijn van Beers

Harry R. Gosker

Daisy J. A. Janssen

Fiona A. H. M. Cleutjens

Frits M. E. Franssen

Martin P. J. van Boxtel

Emiel F. Wouters

Rudolf W. H. M. Ponds

Annemie M. W. J. Schols

*Clinical Nutrition (in press)*

## Abstract

**Background and aims:** Cognitive impairment (CI) and metabolic abnormalities, including the metabolic syndrome (MetS) and sarcopenia, are more prevalent in COPD patients compared to controls without diagnosed lung disease. Because earlier studies have shown these metabolic abnormalities may affect cognitive performance, this study investigated whether cognitive performance is more impaired in subgroups of COPD patients with MetS or sarcopenia.

**Methods:** Cognitive performance patterns of 170 COPD patients referred for pulmonary rehabilitation (53.5% male,  $63.4 \pm 9.4$  years,  $FEV_1$   $54.5 \pm 22.7\%$  predicted) were compared between COPD subgroups stratified by presence of MetS and sarcopenia. Cognitive performance was assessed using a detailed neuropsychological test battery, which measured psychomotor speed (Stroop Color-Word Test, Concept Shifting Test, Letter-Digit Substitution Test), planning (Behavioral Assessment of the Dysexecutive Syndrome), working memory (Visual-Verbal Learning Test, Digit Span), verbal memory (Visual-Verbal Learning Test) and cognitive flexibility (Stroop Color-Word Test, Concept Shifting Test). MetS was determined according to the NCEP ATP-III criteria. Sarcopenia was determined based on decreased appendicular lean mass by dual-energy x-ray absorptiometry and impaired physical performance by 6-minute walking distance.

**Results:** MetS was observed in 54.7% and sarcopenia in 30.0% of COPD patients. The prevalence of general CI was not different between patients with and without MetS (30.4% and 39.0%, respectively) or those with and without sarcopenia (34.0% and 34.5%, respectively, both  $p > 0.05$ ). Domain-specific cognitive performance was not different between metabolic subgroups, but those with sarcopenia displayed a lower prevalence of CI on verbal memory than those without (21.7% and 29.7%, respectively,  $p = 0.011$ ). Only the digit span (working memory) subtest was significantly different between metabolic subgroups, in favor of those without MetS ( $p = 0.017$ ).

**Conclusion:** Cognitive performance was not affected more in COPD patients with sarcopenia or MetS.

## Introduction

Chronic obstructive pulmonary disease (COPD) is often associated with systemic manifestations including metabolic syndrome (MetS) and sarcopenia (1, 2). Cognitive impairment (CI) is also a common, yet often overlooked, comorbidity of COPD, with a recent meta-analysis reporting a 32% prevalence (3). CI in COPD has been associated with lower quality of life (4), a higher need for assistance in aspects of daily living such as medication adherence and self-management (5) and a higher risk of all-cause hospitalization (hazard ratio [HR] for patients with COPD and CI compared to controls without COPD and CI 1.34 [95% confidence interval 1.00-1.80]; HR for patients with COPD but no CI compared to controls without either 1.24 [95% confidence interval 1.06-1.45]) (6), and hospitalized patients with COPD scored significantly worse on a range of cognitive tests compared to their stable counterparts (7). Moreover, patients with CI dropped out of pulmonary rehabilitation (PR) twice as often as those without (23.3 versus 10.3%) (8). Although the etiology of CI in COPD remains largely unknown, it can be speculated that metabolic comorbidities adversely affect cognition in these patients because of shared lifestyle-related risks.

MetS is a cluster of cardiovascular risk factors which, according to a recent review, affects 34% of patients with COPD, a significantly higher proportion than among healthy controls with a comparable body mass index (BMI) (1). MetS is related to reduced levels of multiple indices of cerebral (9, 10) and blood-brain barrier integrity (11) and a reduced blood oxygenation level-dependent (BOLD) imaging signal in frontal and parietal regions (12) in community-dwelling older adults. MetS is also associated with impaired working memory and episodic memory (13), attention and executive functions (13, 14) in healthy elderly, although one study only found significant results in men (13). Some evidence has been found for a dose-response relationship, in which a higher number of MetS symptoms is associated with worse cognitive performance (13). These effects can be long-lasting: in one meta-analysis MetS was associated with a decline in cognitive performance in participants under 70 years old over a follow-up period of up to 16 years (15) and hyperglycemia and insulin resistance have been associated with declines in cognitive performance in a community sample over a 20-year period between ages 53 and 73 (16). The relative contribution of the separate MetS components towards CI remains unclear, although various components or combinations thereof with a significant association with CI have been reported, such as high-density lipoprotein (HDL) cholesterol (17), hyperglycemia (18), waist circumference and hyperglycemia (19), or all components except HDL cholesterol (20).

5

Sarcopenia is defined as a progressive and generalized skeletal muscle disorder that is associated with increased likelihood of adverse outcomes including falls, fractures, physical disability and mortality (21). It occurs in normal ageing but is more prevalent in COPD (21, 22). In healthy elderly populations, a smaller muscle mass has been associated with lower cerebral volume (23) and diminished white matter integrity (24) which lead to poorer cognitive performance. Interestingly, these cerebral abnormalities have also been shown in COPD (25, 26). Furthermore, in healthy elderly, fat-free mass was found to be related to whole-brain volume (27) and total lean mass has been associated with whole-brain volume and white matter volume (28). In a recent review (29) white matter hyperintensity volume and volumes of various brain areas were related to grip strength, but relationships with gait speed were much weaker. Some studies have found a two- to threefold higher risk of developing CI in healthy participants with low muscle mass (30, 31) or functional impairment, defined as a score on the short physical performance battery of less than 9 (32). In one smaller sample ( $N=34$ ) between 71 and 92 years old, the risk was even found to be three- to 4.5-fold, depending on the definition of sarcopenia (33).

The current study therefore aims to investigate whether the presence of MetS or sarcopenia is related to an elevated prevalence and/or different pattern of CI in patients with COPD referred for PR. Investigating this putative connection is important for timely screening of CI in high-risk groups and to optimize and tailor lifestyle interventions.

## Methods

### *Study design*

Cognitive performance of 170 patients with COPD referred for PR was compared between subgroups stratified by sarcopenia or MetS in a cross-sectional design. These patients took part in the COgnitive-PD study (34), which was conducted at CIRO, a center of expertise for patients with chronic organ failure in Horn, The Netherlands, between November 2013 and June 2015. The study was registered in the Dutch Trial Registry (NTR 4215) and approved by the medical ethics committee of Maastricht University Medical Center (NL45127.068.13). The 13 patients who were excluded from the original sample ( $N=183$ ) had missing data on metabolic parameters.

## Patients

Patients with COPD were eligible to take part in this study if they had a diagnosis of COPD based on the Global Initiative for Chronic Obstructive Lung Disease (GOLD) guidelines (35). Those with clinically unstable COPD in the past 4 weeks, a diagnosis of dementia or insufficient mastery of the Dutch language were not eligible to participate. All participants gave written informed consent.

The MAAS population was invited from the Registration Network Family Practices, a patient register of collaborating general practitioners, representative of the Dutch population. Participants were excluded if they suffered from major cerebrovascular pathology, a nervous system tumor or congenital malformation, multiple sclerosis, Parkinsonism, epilepsy, dementia, organic psychosis, schizophrenia, affective psychosis or intellectual disability. Participants with a history of diagnosed lung disease were excluded during the matching process. Demographical (age, sex, educational level), clinical (BMI) and neuropsychological test data was gathered during a 3-hour medical and neurocognitive screening visit. All participants gave written informed consent and both studies were conducted in accordance with the Helsinki Declaration.

## Measurements

### Demographics and lung function

Demographics and lung function (flow volumes and diffusion capacity) of the patients with COPD were assessed during the 3-day assessment at the start of the PR program.

### Metabolic risk profile

MetS and sarcopenia parameters were taken in the COPD population from the 3-day assessment at the start of the PR program. MetS was assessed using the National Cholesterol Education Program Adult Treatment Panel III (NCEP ATP-III) criteria (36), and the sample was divided into groups with and without MetS based on these cutoffs. Sarcopenia was defined as a combination of low muscle mass and functional impairment (21), based on skeletal muscle mass index (SMI) and performance on the 6-minute walking test, respectively. SMI was calculated as appendicular lean mass (ALM) / height<sup>2</sup>. ALM was determined by dual energy x-ray absorptiometry (DEXA). Low muscle mass was defined as an SMI below 7 kg/m<sup>2</sup> for men or below 5.5 kg/m<sup>2</sup> for women, according to Cruz-Jentoft *et al.* (21). Decreased functional performance was defined as a 6-minute walking distance (6MWD) below 80% of the distance predicted using the formula by Troosters *et*



*al.* (37). Participants with low muscle mass and low functional performance based on these cut-offs were assigned to the sarcopenia group; those not meeting both criteria were assigned to the no-sarcopenia group.

## Cognition

Cognitive performance of the patients with COPD was measured using a detailed neuropsychological assessment during a home visit prior to starting PR. See Cleutjens *et al.* (34) for details.

General cognitive performance comprised six test scores: the number of animals named in the animal naming subtest of the Groninger Intelligence Test-II (38), the time to complete card 3 of the Stroop Color-Word Test (39), the time to complete part C of the Concept Shifting Test (40), the number of correct items on the Letter-Digit Substitution Test (41), and the total and delayed recall scores of the Visual-Verbal Learning Test (42).

Performance on five cognitive domains (psychomotor speed, planning, working memory, verbal memory and cognitive flexibility) was assessed using scores on the six tests mentioned above, along with the digit span backwards task (43) and the Behavioral Assessment of the Dysexecutive Syndrome (BADS) (44). See Table 1 for a complete overview.

## *Sample size calculation*

The required sample size was based on a power of 80% and an  $\alpha$  of 0.05. The expected effect size was based on a four-point difference on the St George's Respiratory Questionnaire, and a drop-out rate of 10% was taken into account. These parameters resulted in a required sample size of 183. Additional details can be found in Cleutjens *et al.* (34).

## *Statistical analyses*

Statistical analyses were performed using the Statistical Package for the Social Sciences (SPSS, IBM Corp., Armonk, NY), version 25. A *p*-value below 0.05 was considered statistically significant.

For cognitive profiling, the patients were individually matched in a 1:1 ratio based on age, sex, educational level and BMI to controls from the Maastricht Aging Study (45) (MAAS; approved by the local medical ethics committee [MEC-05-107]). Additional matching on BMI was done because of established independent effects on cognitive performance (46) and because BMI is often affected in MetS

Table 1

*Neuropsychological tests used to determine domain-specific cognitive performance*

Test name	Subscore or subtest	Parameter measures
Psychomotor speed		
SCWT	Card I (time in seconds)	Reading speed
CST	Part A (time in seconds)	Simple motor speed
LDST	Number of correct items in 60 seconds (written and oral)	Psychomotor speed / information processing speed
Planning		
BADS	Key search (profile score: 0-4)	Planning and priority setting
BADS	Zoo map (profile score: 0-4)	
Working memory		
VVLT	Trial 1 (number correct: 0-15)	Immediate verbal span
Digit span	Backward (span: 0-7)	Working memory
Verbal memory		
VVLT	Total recall (number correct: 0-75)	Overall verbal learning and verbal memory after an interval
	Delayed recall (number correct: 0-15)	
	Retention (percentage of words that were retained from the immediate to the delayed score)	
Cognitive flexibility		
SCWT	Card III (time in seconds)	Attentional inhibition of a dominant response
CST	Part C (time in seconds)	Alternating attention

*Adapted from Cleutjens et al. (34). BADS: Behavioral Assessment of the Dysexecutive Syndrome; CST: Concept Shifting Test; LDST: Letter-Digit Substitution Test; SCWT: Stroop Color-Word Test; VVLT: Visual-Verbal Learning Test.*

and sarcopenia. Matching on BMI prevented the effects of BMI-related metabolic risk to confound the effects of MetS- and sarcopenia-related metabolic risk.

Individual raw cognitive test scores of the patients with COPD were converted into Z-scores using the formula  $Z = (X - \mu) / \sigma$ , in which  $X$  is an individual score,  $\mu$  is the mean of the 170 MAAS participants and  $\sigma$  is the standard deviation of the 170 MAAS participants (47). Because the BADS and digit span task were not

administered in MAAS, norm values published elsewhere were used (43, 44). General CI was defined as a Z-score below -1 on two or more of the six contributing tests; domain-specific CI as a Z-score on a specific domain below -1 (47). Domain-specific cognitive performance scores were obtained by averaging the Z-scores of tests belonging to that particular domain.

Within the COPD population, differences on demographic and clinical parameters and cognitive performance were assessed between metabolic subgroups using independent-samples *t*-tests or Mann-Whitney *U*-tests, depending on the distribution of the data. The normality of the distribution was determined using the Kolmogorov-Smirnov test. Differences on categorical variables and prevalence of CI were assessed using  $\chi^2$  tests. The influence of smoking status (current/former/never) and the number of pack years on cognitive performance was assessed using analyses of variance and regression analyses, respectively. Smoking status and the number of pack years served as independent variables, the separate cognitive test scores as dependent variables.

## Results

Demographical characteristics are presented in Table 2. The study group consisted of 170 patients (53.5% male), aged  $63.4 \pm 9.4$  years, with a BMI of  $26.6 \pm 5.6$  kg/m<sup>2</sup> and moderate to severe airflow obstruction ( $FEV_1$   $54.5 \pm 22.7\%$  predicted). General CI was present in 58 patients (34.3%) (see Table 3).

The first trial of the Visual-Verbal Learning Test was the only cognitive test with significant differences between current ( $n=146$ ), former ( $n=22$ ) and never-smokers ( $n=2$ ), due to a significantly higher score by the never-smokers ( $p=0.040$ ). Pack years were only significantly related to the zoo map test of the BADS ( $p=0.008$ ) and the VVLT delayed recall score ( $p=0.046$ ).

### *Metabolic syndrome*

The prevalence of MetS was 54.7%. Patients with MetS had a significantly higher BMI ( $p<0.001$ ) and diffusion capacity ( $p=0.006$ ) than those without. All MetS parameters except for systolic and diastolic blood pressure were significantly different between the groups, all in the expected directions. ALM, SMI, fat mass (all  $p<0.001$ ) and 6MWD % predicted ( $p=0.045$ ) were significantly higher in the group with MetS compared to that without (see Table 2).

General CI was present in 28 COPD patients with MetS and in 30 without (prevalence 30.4% and 39.0%, respectively;  $p=0.245$ ) (see Table 3). Cognitive

Table 2

General clinical and demographical characteristics

Variable	Overall	No MetS	MetS	No sarcopenia	Sarcopenia
General clinical characteristics					
N (male / female) <sup>c</sup>	170 (91/79)	77 (39/38)	93 (52/41)	119 (67/52)	51 (24/27)
Age (years) <sup>b</sup>	63.4±9.4	62.1±10.0	64.5±8.7	63.4±9.6	63.3±9.0
BMI (kg/m <sup>2</sup> ) <sup>a</sup>	26.6±5.6	22.9±4.1	29.7±4.7***	28.8±5.0	21.6±3.5***
Pack years <sup>b</sup>	42.5±24.6	43.0±24.0	42.1±25.2	42.8±24.8	41.9±24.5
6MWD (m) <sup>b</sup>	436.4±109.1	431.2±122.1	440.6±97.6	455.0±104.0	392.9±109.4**
6MWD (% predicted) <sup>b</sup>	68.8±16.2	65.5±17.6	71.5±14.4*	72.4±15.0	60.3±15.9***
Smoking status (current/former/never) <sup>c</sup>	22/146/2	12/65/0	10/81/2	10/109/0	12/37/2**
Lung function parameters					
FEV <sub>1</sub> (% predicted) <sup>a</sup>	54.5±22.7	52.7±26.1	56.0±19.5	60.8±21.2	39.8±19.1***
FVC (% predicted) <sup>a</sup>	100.7±20.8	101.2±24.3	100.2±17.5	104.6±19.0	91.6±22.2***
DLCO (% predicted) <sup>b</sup>	52.4±18.6	48.0±16.0	55.9±19.7**	56.6±18.4	42.4±15.0***
MetS parameters					
Waist circumference (cm) <sup>a</sup>	99.5±16.6	88.4±13.7	108.8±12.7***	104.8±15.0	87.2±13.5***
Fasting glucose (mmol/l) <sup>b</sup>	6.15±1.13	5.76±1.10	6.48±1.06***	6.21±1.03	6.03±1.35*
Triglycerides (mmol/l) <sup>b</sup>	1.50±0.67	1.22±0.45	1.73±0.74***	1.62±0.72	1.21±0.43***
HDL cholesterol (mmol/l) <sup>b</sup>	1.36±0.42	1.47±0.38	1.27±0.42***	1.30±0.41	1.50±0.40**

Table 2 (continued)

Variable	Overall	No MetS	MetS	No sarcopenia	Sarcopenia
Systolic BP (mmHg) <sup>b</sup>	140.5±22.3	134.3±20.1	145.7±22.8**	143.0±22.5	134.8±21.0
Diastolic BP (mmHg) <sup>b</sup>	79.8±12.0	77.9±10.2	81.3±13.2	80.8±11.8	77.3±12.3
Body composition parameters					
FFM (kg) <sup>b</sup>	46.70±10.34	43.24±9.61	49.57±10.08***	49.47±10.10	40.25±7.72***
ALM (kg) <sup>b</sup>	19.85±5.41	18.18±5.08	21.24±5.30***	21.48±5.26	16.05±3.52***
SMI (kg/m <sup>2</sup> ) <sup>a</sup>	6.94±1.41	6.36±1.27	7.41±1.35***	7.47±1.28	5.70±0.80***
Fat mass (kg) <sup>a</sup>	26.16±11.29	18.96±9.07	32.12±9.32***	29.86±10.47	17.53±7.99***
Fat percentage <sup>b</sup>	33.63±9.70	28.55±10.03	37.84±7.06***	35.84±8.82	28.49±9.75***

All data given as mean ± standard deviation, unless noted otherwise. Differences between the groups with and without MetS and those with and without sarcopenia were tested using independent-samples t-tests for variables indicated with <sup>a</sup>, using Mann-Whitney U-tests for those indicated with <sup>b</sup> or using  $\chi^2$  tests for variables indicated with <sup>c</sup>. \*  $p<0.05$ ; \*\*  $p<0.01$ ; \*\*\*  $p<0.001$ .  $\delta$ MWD: 6-minute walking distance; ALM: appendicular lean mass; BMI: body mass index; DLCO: diffusion capacity of the lung for carbon monoxide; FEV<sub>1</sub>: forced expiratory volume in 1 second; FFM: fat-free mass; FVC: forced vital capacity; HDL: high-density lipoprotein; MetS: metabolic syndrome; SMI: skeletal muscle mass index.




Figure 1

Prevalence of domain-specific CI, divided by MetS status

domain scores (see Table 4) and the prevalence of domain-specific CI (see Figure 1) were not different between the groups with and without MetS. The only significant difference between the groups on the separate cognitive subtests was a better performance on the digit span backwards test by the group without MetS ( $p=0.017$ ) (see Table 4).

### Sarcopenia

The prevalence of sarcopenia was 30.0%. Patients with sarcopenia had a significantly lower BMI ( $p<0.001$ ) and 6MWD than those without, expressed as




Figure 2

Prevalence of domain-specific CI, divided by sarcopenia status. \*  $p < 0.05$ .

absolute values ( $p = 0.001$ ) and percentage predicted ( $p < 0.001$ ). The distribution of current, former and never-smokers was significantly different between the groups, with a significantly higher proportion of current and never-smokers in the group with sarcopenia ( $p = 0.002$ ). All lung function parameters were significantly higher in the group without sarcopenia, indicating better lung function in the patients without sarcopenia (all  $p < 0.001$ ), and all MetS and body composition parameters except systolic and diastolic blood pressure were significantly different between the groups (see Table 2).

Table 3

Cognitive test scores used to determine general CI

Variable	Overall (N=170)	No MetS (n=77)	MetS (n=93)	No sarcopenia (n=119)	Sarcopenia (n=51)
Prevalence of general CI (%) <sup>c</sup>	34.3	39.0	30.4	34.5	34.0
GIT verbal fluency (# of animal names) <sup>b</sup>	22.0±6.7	21.6±6.9	22.3±6.5	22.1±6.6	21.6±7.0
LDST 60 sec oral (# correct) <sup>b</sup>	33.7±7.5	34.3±8.6	33.2±6.5	33.4±7.5	34.4±7.5
WLT total recall trial 1-5 (# correct) <sup>a</sup>	41.6±11.0	42.6±11.1	40.8±10.9	41.8±11.4	41.3±10.0
WLT delayed recall (# correct) <sup>b</sup>	7.8±3.6	8.1±3.7	7.6±3.5	7.7±3.8	8.2±3.0
SCWT, card 3 (s) <sup>b</sup>	54.9±21.4 †††	54.7±22.2	55.0±20.9	55.5±23.9	53.4±14.3
CST-C (s) <sup>b</sup>	44.6±22.6	44.9±27.4	44.4±17.9	45.2±23.2	43.3±21.4

All data given as mean ± standard deviation, unless noted otherwise. Differences between the groups with and without MetS and those with and without sarcopenia were tested using independent-samples t-tests for variables indicated with <sup>a</sup>, using Mann-Whitney U-tests for those indicated with <sup>b</sup> or using  $\chi^2$  tests for variables indicated with <sup>c</sup>. †††  $p<0.001$  compared to the MAAS population. CST: Concept Shifting Test; GIT: Groninger Intelligence Test; LDST: Letter-Digit Substitution Test; MetS: metabolic syndrome; SCWT: Stroop Color-Word Test; WLT: Visual-Verbal Learning Test.



Table 4

Cognitive test scores used to determine domain-specific CI

Variable	Overall (N=170)	No MetS (n=77)	MetS (n=93)	No sarcopenia (n=119)	Sarcopenia (n=51)
<i>Psychomotor speed (Z-score)<sup>b</sup></i>	-0.02±1.12	-0.01±1.44	-0.02±0.77	-0.02±1.16	-0.01±1.04
SCWT, card 1 (s) <sup>b</sup>	17.9±4.5	18.1±5.6	17.8±3.3	18.0±5.0	17.9±3.1
CST-A (s) <sup>b</sup>	25.4±12.6	26.6±17.0	24.4±7.1	25.2±13.3	25.9±11.1
LDST 60 sec read (# correct) <sup>b</sup>	27.4±7.4	28.4±8.2	26.6±6.5	27.4±7.2	27.5±7.8
LDST 60 sec oral (# correct) <sup>b</sup>	33.7±7.5	34.3±8.6	33.2±6.5	33.4±7.5	34.4±7.5
<i>Planning (Z-score)<sup>b</sup></i>	-0.05±0.68	-0.04±0.67	-0.05±0.68	-0.05±0.69	-0.05±0.66
BADS key search profile score <sup>b</sup>	2.5±1.4	2.6±1.4	2.5±1.4	2.5±1.4	2.6±1.4
BADS zoo map profile score <sup>b</sup>	2.4±0.8	2.4±0.8	2.4±0.8	2.4±0.8	2.3±0.8
<i>Working memory (Z-score)<sup>b</sup></i>	-0.51±0.77	-0.45±0.82	-0.56±0.73	-0.52±0.80	-0.48±0.72
WLT trial 1 (# correct) <sup>b</sup>	4.7±1.9	4.6±1.9	4.7±1.9	4.8±1.9	4.5±1.8
Digit span backwards (# correct) <sup>b</sup>	3.0±1.1	3.2±1.1	2.9±1.0 *	2.9±1.1	3.3±0.9
<i>Verbal memory (Z-score)<sup>a</sup></i>	-0.11±1.02	-0.04±1.09	-0.16±0.97	-0.15±1.08	-0.01±0.86
WLT total recall trial 1-5 (# correct) <sup>a</sup>	41.6±11.0	42.6±11.1	40.8±10.9	41.8±11.4	41.3±10.0
WLT delayed recall (# correct) <sup>b</sup>	7.8±3.6	8.1±3.7	7.6±3.5	7.7±3.8	8.2±3.0
WLT retention score (%) <sup>b</sup>	72.8±25.7	72.7±28.1	72.8±23.6	70.7±26.6	77.5±23.0
<i>Cognitive flexibility (Z-score)<sup>b</sup></i>	-0.51±1.39	-0.52±1.59	-0.51±1.22	-0.56±1.51	-0.41±1.09
SCWT, card 3 (s) <sup>b</sup>	54.9±21.4	54.7±22.2	55.0±20.9	55.5±23.9	53.4±14.3

Table 4 (continued)

Variable	Overall (N=170)	No MetS (n=77)	MetS (N=93)	No sarcopenia (n=119)	Sarcopenia (n=51)
CST-C (s) <sup>b</sup>	44.6±22.6	44.9±27.4	44.4±17.9	45.2±23.2	43.3±21.4

All data given as mean ± standard deviation, unless noted otherwise. Differences between the groups with and without MetS and those with and without sarcopenia were tested using independent-samples t-tests for variables indicated with <sup>a</sup> or Mann-Whitney U-tests for those indicated with <sup>b</sup>. \* p<0.05. BADS: Behavioral Assessment of the Dysexecutive Syndrome; CST: Concept Shifting Test; LDST: Letter-Digit Substitution Test; MetS: metabolic syndrome; SCWT: Stroop Color-Word Test; VLT: Visual-Verbal Learning Test.

General CI was present in 17 COPD patients with sarcopenia and in 41 without (prevalence 34.0% and 34.5%, respectively;  $p=0.955$ ) (see Table 3). The prevalence of CI on the verbal memory domain was significantly lower in those with sarcopenia compared to those without ( $p=0.011$ ) (see Figure 2), but no cognitive domain scores or separate cognitive tests were significantly different between the groups (see Table 4).

## Discussion

The current study shows a higher prevalence of MetS and sarcopenia in patients with COPD eligible for PR, than reported earlier in patients with COPD both in and outside of a PR context (1, 48, 49). The prevalence of CI was comparable to that in a recent systematic review (3). There was no different pattern of cognitive performance in the COPD subgroups with MetS or sarcopenia compared to the subgroups without.

MetS and sarcopenia and their separate parameters did not significantly affect cognitive performance in our study group. This does not exclude the possibility that common and shared lifestyle determinants of metabolic and cognitive disturbances in COPD, including physical activity (50, 51) and a low-quality diet (52, 53), may explain the high prevalence of impaired cognitive performance. Although smoking can also affect cognitive performance (54, 55), its influence in the current study was limited. Further research is warranted to disentangle the relative contribution of these modifiable lifestyle factors in the etiology and progression of CI in COPD.

Our results are at odds with earlier research that did associate MetS (56-58) and sarcopenia (59, 60) with poorer cognitive performance in participants without COPD. Other studies found non-significant (61-63) or mixed results (17, 18). These inconsistent results can result from different sources, such as differences in age ranges, in the operationalization of cognitive performance and CI, in the operationalization of MetS or sarcopenia or from lack of sufficient power. Secondly, methodological weaknesses in cognitive screening can distort results. Many previous studies gauged cognition using a single screening tool, which is less sensitive than a comprehensive test battery, or excluded participants with pre-existing CI (e.g., Mini-Mental State Examination score  $\leq 24$ ). The use of a screening tool and selectively excluding certain participant groups both reduce variance on cognitive outcome measures and could lead to underestimating effect sizes. Thirdly, not all of the MetS components are equally strongly associated

with cognition (16-18). Also in our sample, only glucose levels were significantly positively related to cognition (data not shown).

Strengths of the current study include the use of a comprehensive neuropsychological test battery, rather than a single screening tool, to investigate cognitive performance. The calculation of Z-scores based on case-control matching allowed us to reliably compare cognitive performance of the patient population to controls corrected for known determinants of cognitive functioning.

A strength but also potential limitation includes the fact that our study population consisted of patients in tertiary care. The prevalence of metabolic derangements is high in this population, but many patients have complex pathology and comorbidities, which might limit the generalizability of the results to COPD per se. Secondly, we were unable to control for confounding factors such as socio-economic status or COPD disease duration, following the studies by Cleutjens *et al.* (47) and many others. Thirdly, like most other comparable studies, we were constrained to a cross-sectional study design, which limits our ability to draw conclusions about causality.

In conclusion, COPD subgroups with MetS and sarcopenia exhibited comparable patterns of cognitive performance and impairment relative to those without. Because earlier research has shown poorer cognitive performance in patients with COPD compared to healthy controls, COPD constitutes an indication for cognitive screening, but the presence of MetS or sarcopenia in patients with COPD does not.

## References

1. Cebzon Lipovec N, Beijers RJ, Van den Borst B, Doehner W, Lainscak M, Schols AMWJ. The prevalence of metabolic syndrome in chronic obstructive pulmonary disease: a systematic review. *COPD*. 2016;13(3):399-406.
2. Benz E, Trajanoska K, Lahousse L, Schoufour JD, Terzikhan N, De Roos E, et al. Sarcopenia in COPD: a systematic review and meta-analysis. *European Respiratory Review*. 2019;28:190049.
3. Yohannes AM, Chen W, Moga AM, Leroi I, Connolly MJ. Cognitive impairment in chronic obstructive pulmonary disease and chronic heart failure: a systematic review and meta-analysis of observational studies. *JAMDA*. 2017;18:451.e1-e11.
4. Roncero C, Campuzano AI, Quintano JA, Molina J, Perez J, Miravittles M. Cognitive status among patients with chronic obstructive pulmonary disease. *International Journal of COPD*. 2016;11:543-51.
5. Baird C, Lovell J, Johnson M, Shiell K, Ibrahim JE. The impact of cognitive impairment on self-management in chronic obstructive pulmonary disease: a systematic review. *Respiratory Medicine*. 2017;129:130-9.
6. Chang SS, Chen S, McAvay GJ, Tinetti ME. Effect of coexisting chronic obstructive pulmonary disease and cognitive impairment on health outcomes in older adults. *Journal of the American Geriatric Society*. 2012;60(10):1839-46.
7. Dodd JW, Charlton RA, Van den Broek MD, Jones PW. Cognitive dysfunction in patients hospitalized with acute exacerbation of COPD. *Chest*. 2013;144(1):119-27.
8. Cleutjens FAHM, Spruit MA, Ponds RWHM, Vanfleteren LEGW, Franssen FM, Dijkstra JB, et al. The impact of cognitive impairment on efficacy of pulmonary rehabilitation in patients with COPD. *JAMDA*. 2017;18(5):420-6.
9. Sala M, De Roos A, Van den Berg A, Altmann-Schneider I, Slagboom PE, Westendorp RG, et al. Microstructural brain tissue damage in metabolic syndrome. *Diabetes Care*. 2014;37(2):493-500.
10. Alfaro FJ, Gavrieli A, Saade-Lemus P, Lioutas V-A, Upadhyay J, Novak V. White matter microstructure and cognitive decline in metabolic syndrome: a review of diffusion tensor imaging. *Metabolism*. 2018;78:52-68.
11. Mauro C, De Rosa V, Marelli-Berg F, Solito E. Metabolic syndrome and the immunological affair with the blood-brain barrier. *Frontiers in Immunology*. 2015;2015(5):677.
12. Shigaeff N, Amaro E, Franco FGM, Jacinto AF, Chiochetta G, Cendoroglo MS, et al. Functional magnetic resonance imaging response as an early biomarker of cognitive decline in elderly patients with metabolic syndrome. *Archives of Gerontology and Geriatrics*. 2017;73:1-7.
13. Cavalieri M, Ropele S, Petrovic K, Pluta-Fuerst A, Homayoon N, Enzinger C, et al. Metabolic syndrome, brain magnetic resonance imaging, and cognition. *Diabetes Care*. 2010;33(12):2489-95.

14. Falkowski J, Atchison T, DeButte-Smith M, Weiner MF, O'Bryant S. Executive functioning and the metabolic syndrome: a Project FRONTIER study. *Archives of Clinical Neuropsychology*. 2014;29:47-53.
15. Siervo M, Harrison SL, Jagger C, Robinson L, Stephan BCM. Metabolic syndrome and longitudinal changes in cognitive function: a systematic review and meta-analysis. *Journal of Alzheimer's Disease*. 2014;41:151-61.
16. Tortelli R, Lozupone M, Guerra V, Barulli MR, Imbimbo BP, Capozzo R, *et al.* Midlife metabolic profile and the risk of late-life cognitive decline. *Journal of Alzheimer's Disease*. 2017;59(1):121-30.
17. Feinkohl I, Janke J, Hadzidiakos D, Slooter A, Winterer G, Spies C, *et al.* Associations of the metabolic syndrome and its components with cognitive impairment in older adults. *BMC Geriatrics*. 2019;19:77.
18. Overman MJ, Pendleton N, O'Neill TW, Bartfai G, Casanueva FF, Forti G, *et al.* Glycemia but not the metabolic syndrome is associated with cognitive decline: findings from the European Male Ageing Study. *American Journal of Geriatric Psychiatry*. 2017;25(6):662-71.
19. Lai MMY, Ames DJ, Cox KL, Ellis KA, Sharman MJ, Hepworth G, *et al.* Association between cognitive function and clustered cardiovascular risk of metabolic syndrome in older adults at risk of cognitive decline. *Journal of Nutrition, Health and Aging*. 2020;24(3):300-4.
20. Chen L, Hong W, Yang H, Dong S, Peng Z, Zhou H. Effects of metabolic syndrome on cognitive impairment with white matter lesions. *European Neurology*. 2019;82(4-6):99-105.
21. Cruz-Jentoft AJ, Bahat G, Bauer JM, Boirie Y, Bruyère O, Cederholm T, *et al.* Sarcopenia: revised European consensus on definition and diagnosis. *Age and Ageing*. 2019;48(1):16-31.
22. Trajanoska K, Schoufour JD, Darweesh SKL, Benz E, Medina-Gomez C, Alferink LJM, *et al.* Sarcopenia and its clinical correlates in the general population: the Rotterdam Study. *Journal of Bone and Mineral Research*. 2018;33(7):1209-18.
23. Kilgour AHM, Ferguson KJ, Gray CD, Deary IJ, Wardlaw JM, MacLulich AMJ, *et al.* Neck muscle cross-sectional area, brain volume and cognition in healthy older men: a cohort study. *BMC Geriatrics*. 2013;13:20.
24. Kohara K, Okada Y, Ochi M, Ohara M, Nagai T, Tabara Y, *et al.* Muscle mass decline, arterial stiffness, white matter hyperintensity, and cognitive impairment: Japan Shimanami Health Promoting Program study. *Journal of Cachexia, Sarcopenia and Muscle*. 2017;8(4):557-66.
25. Li J, Fei GH. The unique alterations of hippocampus and cognitive impairment in chronic obstructive pulmonary disease. *Respiratory Research*. 2013;14:140.

26. Spilling CA, Jones PW, Dodd JW, Barrick TR. White matter lesions characterise brain involvement in moderate to severe chronic obstructive pulmonary disease, but cerebral atrophy does not. *BMC Pulmonary Medicine*. 2017;17:92.
27. Heymsfield SB, Müller MJ, Bosy-Westphal A, Thomas D, Shen W. Human brain mass: similar body composition associations as observed across mammals. *American Journal of Human Biology*. 2012;24:479-85.
28. Burns JM, Johnson DK, Watts A, Swerdlow RH, Brooks WM. Reduced lean mass in early Alzheimer disease and its association with brain atrophy. *Archives of Neurology*. 2010;67(4):428-33.
29. Kilgour AHM, Todd OM, Starr JM. A systematic review of the evidence that brain structure is related to muscle structure and their relationship to brain and muscle function in humans over the lifecourse. *BMC Geriatrics*. 2014;14:85.
30. Peng T-C, Chen W-L, Wu L-W, Chang Y-W, Kao T-W. Sarcopenia and cognitive impairment: a systematic review and meta-analysis. *Clinical Nutrition*. 2019;S0261-5614(19):33200-5.
31. Cabett Cipolli G, Sandes Yassuda M, Aprahamian I. Sarcopenia is associated with cognitive impairment in older adults: a systematic review and meta-analysis. *Journal of Nutrition, Health and Aging*. 2019;23:525-31.
32. Moon JH, Moon JH, Kim KM, Choi SH, Lim S, Park KS, *et al*. Sarcopenia as a predictor of future cognitive impairment in older adults. *Journal of Nutrition, Health and Aging*. 2016;20(5):496-502.
33. Papachristou E, Ramsay SE, Lennon LT, Papacosta O, Iliffe S, Whincup PH, *et al*. The relationships between body composition characteristics and cognitive functioning in a population-based sample of older British men. *BMC Geriatrics*. 2015;15:172.
34. Cleutjens FAHM, Wouters EFM, Dijkstra JB, Spruit MA, Franssen F, Vanfleteren LEGW, *et al*. The COgnitive-Pulmonary Disease (COgnitive-PD) study: protocol of a longitudinal observational comparative study on neuropsychological functioning of patients with COPD. *BMJ Open*. 2014;4(3):e004495.
35. Vestbo J, Hurd SS, Agustí AG, Jones PW, Vogelmeier C, Anzueto A, *et al*. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *American Journal of Respiratory and Critical Care Medicine*. 2013;187(4):347-65.
36. Grundy SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, *et al*. Diagnosis and management of the metabolic syndrome: an American Heart Association / National Heart, Lung and Blood Institute scientific statement. *Circulation*. 2005;112(17):2735-52.
37. Troosters T, Gosselink R, Decramer M. Six minute walking distance in healthy elderly subjects. *European Respiratory Journal*. 1999;14(2):270-4.

38. Barelids DPH. Groninger Intelligentie Test 2 (GIT-2): handleiding. Amsterdam, The Netherlands: Harcourt Publishers; 2004.
39. Van der Elst W, Van Boxtel MP, Van Breukelen GJ, Jolles J. The Stroop Color-Word Test: influence of age, sex and education; and normative data for a large sample across the adult age range. *Assessment*. 2006;13(1):62-79.
40. Van der Elst W, Van Boxtel MP, Van Breukelen GJ, Jolles J. The Concept Shifting Test: adult normative data. *Psychological Assessment*. 2006;18(4):424-32.
41. Van der Elst W, Van Boxtel MP, Van Breukelen GJ, Jolles J. The Letter Digit Substitution Test: normative data for 1,858 healthy participants aged 24-81 from the Maastricht Aging Study (MAAS): influence of age, education and sex. *Journal of Clinical and Experimental Neuropsychology*. 2006;28(6):998-1009.
42. Van der Elst W, Van Boxtel MP, Van Breukelen GJ, Jolles J. Rey's Verbal Learning Test: normative data for 1855 healthy participants aged 24-81 years and the influence of age, sex, education, and mode of presentation. *Journal of the International Neuropsychological Society*. 2005;11(3):290-302.
43. Monaco M, Costa A, Caltagirone C, Carlesimo GA. Forward and backward span for verbal and visuo-spatial data: standardization and normative data from an Italian adult population. *Neurological Sciences*. 2013;34(5):749-54.
44. Wilson BA, Alderman N, Burgess PW, Emslie HE, Evans JJ. Behavioral Assessment of the Dysexecutive Syndrome. Bury St Edmunds, UK: Thames Valley Test Company; 1996.
45. Jolles J, Van Boxtel MPJ, Ponds RW, Metsemakers JFM, Houx PJ. The Maastricht Aging Study (MAAS): The longitudinal perspective of cognitive aging. *Tijdschrift voor Gerontologie en Geriatrie*. 1998;29(3):120-9.
46. Arvanitakis Z, Capuano AW, Bennett DA, Barnes LL. Body mass index and decline in cognitive function in older black and white persons. *Journal of Gerontology A*. 2018;73(2):198-203.
47. Cleutjens FAHM, Franssen F, Spruit MA, Vanfleteren LEGW, Gijssen C, Dijkstra JB, et al. Domain-specific cognitive impairment in patients with COPD and control subjects. *International Journal of Chronic Obstructive Pulmonary Disease*. 2017;12:1-11.
48. Byun MK, Cho EN, Chang J, Ahn CM, Kim HJ. Sarcopenia correlates with systemic inflammation in COPD. *International Journal of Chronic Obstructive Pulmonary Disease*. 2017;12:669-75.
49. Jones SE, Maddocks M, Kon SSC, Canavan JL, Nolan CM, Clark AL, et al. Sarcopenia in COPD: prevalence, clinical correlates and response to pulmonary rehabilitation. *Thorax*. 2015;70:213-8.
50. Kantorowski A, Wan ES, Homsy D, Kadri R, Richardson CR, Moy ML. Determinants and outcomes of change in physical activity in COPD. *ERJ Open Research*. 2018;4(3):00054-2018.



51. Blondell SJ, Hammersley-Mather R, Veerman JL. Does physical activity prevent cognitive decline and dementia? A systematic review and meta-analysis of longitudinal studies. *BMC Public Health*. 2014;14:510.
52. Van de Boel C, Mattijssen-Verdonschot C, Van Melick PPMJ, Spruit MA, Franssen FME, Wouters EFM, *et al*. Quality of dietary intake in relation to body composition in patients with chronic obstructive pulmonary disease eligible for pulmonary rehabilitation. *European Journal of Clinical Nutrition*. 2014;68(2):159-65.
53. Abbatecola AM, Russo M, Barbieri M. Dietary patterns and cognition in older persons. *Current Opinion in Clinical Nutrition and Metabolic Care*. 2018;21(1):10-3.
54. Swan GE, Leslov-Schlaggar CN. The effects of tobacco smoke and nicotine on cognition and the brain. *Neuropsychological Review*. 2007;17:259-73.
55. Waisman Campos M, Serebrisky D, Castaldelli-Maia JM. Smoking and cognition. *Current Drug Abuse Reviews*. 2016;9:1-4.
56. Lee EY, Lee SJ, Kim KM, Yun YM, Song BM, Kim JE, *et al*. Association of metabolic syndrome and 25-hydroxyvitamin D with cognitive impairment among elderly Koreans. *Geriatrics & Gerontology International*. 2017;17(7):1069-75.
57. Viscogliosi G, Chiriac IM, Andreozzi P, Ettore E. The metabolic syndrome predicts longitudinal changes in clock drawing test performance in older nondemented hypertensive individuals. *American Journal of Geriatric Psychiatry*. 2016;24(5):359-63.
58. Viscogliosi G, Donfrancesco C, Palmieri L, Giampaoli S. The metabolic syndrome and 10-year cognitive and functional decline in very old men - a population-based study. *Archives of Gerontology and Geriatrics*. 2017;70:62-6.
59. Tolea MI, Galvin JE. Sarcopenia and impairment in cognitive and physical performance. *Clinical Interventions in Aging*. 2015;10:663-71.
60. Tolea MI, Chrisphonte S, Galvin JE. Sarcopenic obesity and cognitive performance. *Clinical Interventions in Aging*. 2018;13:1111-9.
61. Martinez-Miller EE, Kohl III HW, Barlow CE, Willis BL, DeFina LF. Metabolic syndrome and cognitive impairment among high socioeconomic, nondemented older US adults. *Journal of the American Geriatric Society*. 2019;67(7):1437-43.
62. Auyeung TW, Lee JSW, Kwok T, Woo J. Physical frailty predicts future cognitive decline - a four-year prospective study in 2737 cognitively normal older adults. *Journal of Nutrition, Health and Aging*. 2011;15(8):690-4.
63. Lasaitė L, Krasauskienė A. Psychological state, quality of life and body composition in postmenopausal women with osteoporosis in Lithuania. *Archives of Osteoporosis*. 2009;4(1-2):85-90.





# Chapter 6

Efficacy of working memory training in  
chronic obstructive pulmonary disease: the  
randomized placebo-controlled Cogtrain  
trial

Martijn van Beers

Sarah W. Mount

Katrijn Houben

Harry R. Gosker

Lisanne Schuurman

Frits M. E. Franssen

Daisy J. A. Jansen

Annemie M. W. J. Schols

*Submitted*

# **Chapter 7**

## General Discussion

Leading a healthy lifestyle is important for patients with COPD. It benefits their overall health status, for instance through slowing down disease progression. Nevertheless, the lifestyle of many patients with COPD is unhealthy: some of them still smoke, they are often physically inactive (1, 2) and consume an unhealthy diet (3-5). This thesis reported that patients' motivational patterns are associated with their physical activity and dietary quality, which tentatively indicates that targeted motivational counseling might be effective in order to achieve or consolidate a healthy lifestyle. However, existing evidence also indicated that improving cognitive impairment (CI) or cognitive functioning could be effective to this end. Unfortunately, CI remains an under-recognized manifestation of COPD and our working memory training (WMT) intervention could not improve overall cognitive functioning or adherence to a healthy lifestyle.

This chapter will discuss how the clinical recognition and assessment of CI in COPD could be improved, and it will discuss various other potentially effective strategies to improve cognitive functioning in patients with COPD and their adherence to a healthy lifestyle.

## **Cognitive impairment as under-recognized COPD manifestation**

CI is highly prevalent in COPD, with a prevalence of around one in three (6), and cognitive performance substantially influences patients' daily lives. Memory, cognitive flexibility and visual processing are related to self-management skills, such as the ability to adhere to one's medication regime (7, 8), and CI in general is associated with increased risk for exacerbations and hospitalizations (9-11). Nevertheless, many patients and health care professionals under-recognize CI as a comorbidity of COPD (12). Even the GOLD guidelines (13), outlining the standards in COPD management, do not address cognitive functioning and its influence on disease management and daily-life outcomes.

Cognitive screening could be a valuable addition to COPD management, as it could enable timely intervention and prevention of the harmful consequences of CI. Cognitive screening is not time- or labor-intensive: instruments such as the Mini-Mental State Examination (MMSE) (14), the Montreal Cognitive Assessment (MoCA) (15) or the Addenbrooke's Cognitive Examination-Revised (ACE-R) (16) take less than ten minutes to administer and are sensitive to detect CI in patients with COPD. They are easy to score and to interpret because of clear cut-off

values for CI. Moreover, patients with COPD themselves do not regard cognitive screening as a strange addition to standard diagnostics (12).

Nurses or lab technicians could periodically conduct this cognitive screening, for instance in conjunction with patients' routine lung function evaluation, which usually takes place biannually in The Netherlands. Cognitive screening could be followed up in a similar way as spirometric evaluation: if it reveals impairment or decline, the general practitioner or chest physician can be informed and advised to refer the patient for more comprehensive investigation. Cognitive abnormalities could lead, for example, to referral to a memory clinic in the hospital, which can administer comprehensive neuropsychological assessment.

### *Metabolic syndrome and sarcopenia as potential indications for screening*

Multifactorial involvement of disease-specific determinants, comorbidities and unhealthy lifestyle behaviors in the emergence of CI in COPD makes some of these factors potentially useful to target screening at specific at-risk subgroups of patients with COPD. Metabolic abnormalities such as the metabolic syndrome (MetS) and sarcopenia, for instance, are common conditions in patients with COPD compared to healthy controls (17, 18) and have been related to the prevalence of CI in non-COPD populations (19-23). However, our research showed no independent contribution of MetS and sarcopenia towards CI in patients with COPD who were referred for pulmonary rehabilitation (PR). This could be specific for this COPD subgroup and different in population-based studies. Future research is furthermore indicated to identify the potential of other comorbidities or lifestyle factors in the risk screening for CI in COPD.

Alternatively, COPD may have a much larger effect on cognitive performance than MetS and sarcopenia. The impact of COPD on cognitive functioning might therefore overshadow the (perhaps negligible) impact of MetS and sarcopenia. This conjecture is supported by the relatively high prevalence of CI in the patients who were investigated in our research (almost 40 percent), regardless of MetS or sarcopenia status. This prevalence was determined by comparing the patients' cognitive performance to that of non-COPD controls. COPD-general effects on cognitive performance could be biologically mediated by accelerated ageing. Accelerated biological ageing is a hallmark of COPD (24) and has also been related to poorer cognitive performance (25-27) and accelerated cognitive decline (26, 28, 29) in otherwise healthy participants.

## Improving cognitive performance

In case CI is diagnosed, it should ideally be treated, especially in light of the serious consequences of CI in patients with COPD. We chose to administer WMT as the role of WM underlying general cognition (30) made it a promising target to improve overall cognitive performance. However, the results of the Cogtrain trial as well as earlier WMT interventions were disappointing. In the Cogtrain trial as well as earlier studies, WMT appeared to be able to improve cognitive performance on the initially trained tasks, but near transfer (i.e., towards comparable cognitive tasks) was scarce and far transfer (i.e., towards unrelated cognitive tasks) was virtually non-existent (31-33). To date only one study investigated another type of cognitive training in patients with COPD. This intervention, targeting attention and logical-deductive thinking with a lower training load than the Cogtrain trial, was also ineffective (34).

Based on this evidence, WMT cannot be recommended for use in future studies or clinical practice, at least not as a stand-alone intervention in its current form.

*Targeting cognitively well- or poorly-performing patients: who benefits?*

Baseline cognitive performance in the Cogtrain trial was highly variable. The median ACE-R score of 88 was relatively poor, but the highest-performing quartile of patients showed normal cognitive performance with scores above 94. This variability could also explain the non-significant results of the trial, as WMT might have different effects on patients with poor and good baseline cognitive performance. This has not been investigated in the Cogtrain trial, as the subgroups resulting from its relatively small sample size would be too small to draw any meaningful conclusions.

On the one hand, cognitively well-performing patients might have diluted effects that were present in more poorly-performing ones. Transfer from improved performance on one cognitive task towards another is unlikely to occur if participants do not improve on the trained tasks (35), and the available room for improvement is largest in cognitively poor patients. On the other hand, poorly performing patients might also have diluted effects that were present in the better ones. Plasticity is a prerequisite for learning and transfer, and it could be argued that the potential for cognitive plasticity (i.e., the ability to learn and use multiple new cognitive strategies) and neural plasticity (i.e., cerebral physiological adaptability) is likely largest in those with better initial cognitive performance.



### *Daily-life cognitive stimulation*

Improving cognitive functioning does not require standardized or formalized cognitive training. Cognitive stimulation associated with daily-life activities such as playing chess (36), completing sudokus (37) or even the degree of social contact (38, 39) are also associated with improved cognitive functioning. These activities have multiple advantages over formalized cognitive training: their structure is less rigid and they have a much more 'gamified' character, so patients will likely be more willing to engage in them and adhere to them. They are easy to incorporate into daily life. This increases the likelihood that these activities become a habit, which makes it easier to adhere to them in the longer term.

### *Physical activity, exercise training and the muscle-brain axis*

Physical activity and exercise training are among the most effective ways to improve brain health and cognitive functioning (40-42). These effects are mediated among others by increased cardiac output, social interaction in the case of group exercise (43) and extensive muscle-brain crosstalk.

This crosstalk takes place among others through the effects of exercise-induced release of myokines such as cathepsin B or FNDC5 (44). These and other factors mediate the upregulation of neurotransmitters such as brain-derived neurotrophic factor (BDNF), vascular endothelial growth factor (VEGF) and insulin growth factor-1 (IGF-1) (44-46). These neurotransmitters improve brain health by facilitating neurogenesis, neuronal survival, differentiation, myelination and memory formation (43, 47). A large part of these effects takes place in the hippocampus (44, 48), which is the primary brain area responsible for memory formation.

The kynurenine pathway also plays a key role in this muscle-brain crosstalk. This pathway mainly converts tryptophan into kynurenine and further into quinolinic acid (an N-methyl-D-aspartate receptor [NMDAR] agonist which induces excitotoxicity in the brain) or kynurenic acid (an NMDAR antagonist with neuroprotective properties) (49, 50). Quinolinic and kynurenic acid cannot cross the blood-brain barrier, but kynurenine can (51). Kynurenine should therefore ideally be metabolized in the periphery, to prevent it from crossing the blood-brain barrier and being metabolized into neurotoxic compounds in the central nervous system. This risk is especially present in patients with COPD, because inflammation, which is systemically heightened in patients with COPD (13), promotes the conversion of kynurenine into quinolinic acid instead of kynurenic acid (50). Accordingly, the

kynurenine-to-tryptophan ratio (KTR) has been negatively related to cognitive performance (52).

Exercise promotes peripheral kynurenine metabolism (and therefore potentially benefits cognitive performance) through increasing PGC-1 $\alpha$  and PPAR $\alpha/\delta$  expression in the skeletal muscle (53). This leads to increased skeletal muscle kynurenine aminotransferase (KAT) expression (49, 53), which in turn promotes the conversion of kynurenine into kynurenic acid (53). Indeed, the peripheral KTR of healthy young adults was significantly lower 1 hour after exercise compared to baseline (49). Indirectly improving cognitive performance in COPD via such exercise-induced muscular improvements makes sense in theory, as their plasma kynurenine levels are significantly higher than those of healthy controls and their skeletal muscle KAT expression is significantly lower (54).

### *Nutritional supplementation, dietary management and the gut-brain axis*

Brain health and cognitive performance may also be improved by targeting the gut microbiome. The gut microbiome communicates with the brain through the 'gut-brain axis' (55). This axis consists of many different pathways. It releases pro- and anti-inflammatory cytokines such as interleukin-1 and -6, which modulate the activity of the hypothalamic-pituitary-adrenal gland (HPA) axis (55). It secretes short-chain fatty acids which activate microglial cells that have anti-inflammatory effects in the brain (56). It modulates the activity of the vagus nerve, which is associated with improved cognitive functioning in the general population when it is done exogenously (57, 58), and it also regulates the levels of (precursors of) important neurotransmitters such as serotonin, gamma aminobutyric acid (GABA), noradrenalin, dopamine and acetylcholine (55, 56).

The composition of the gut microbiome and the functioning of the gut-brain axis can be improved by administering so-called 'psychobiotics'. Psychobiotics are "pre- and probiotics that, ingested in appropriate quantities, have a positive mental health benefit" (55). Fructooligosaccharides (FOS) and galactooligosaccharides (GOS), for instance, stimulate the growth of intrinsic commensal microbiota, which are bacteria in the gut that positively influence the diversity of the gut microbiome (55). The current evidence for the beneficial effects of psychobiotics is strongest for stress, anxiety and depression (59); surprisingly, there is a paucity of research into the effects of psychobiotic supplementation on cognitive performance. Future studies may therefore further investigate this topic, and whether and how psychobiotic supplementation is related to the composition and diversity of the gut microbiome.

Various nutrients can also modulate the gut microbiome and gut-brain axis. The NUTRECOVER trial, which is currently ongoing within the department of respiratory medicine at Maastricht University, will unravel the effects of a nutritional supplement containing tryptophan, vitamin D, omega-3 polyunsaturated fatty acids (PUFA) and prebiotic fibers on among others cognitive performance in patients with COPD who have been hospitalized with an acute exacerbation. This 'cocktail' may simultaneously moderate the kynurenine pathway and the gut-brain axis. The anti-inflammatory properties of PUFAs, such as eicosapentaenoic acid (EPA) or docosahexaenoic acid (DHA) (60), could shift the metabolism of the supplemented tryptophan and subsequently kynurenine in the brain towards kynurenic acid instead of quinolinic acid. Additionally, the prebiotic fibers may improve the composition of the gut microbiome (59), which might in turn also benefit the kynurenine pathway (61). Clinical trials have already shown positive effects of PUFA and dietary fiber supplementation on cognitive performance in cognitively healthy elderly and those with mild CI (62-64). A lower vitamin D status has been associated with poorer cognitive performance in (older) adults without dementia, although the benefits of supplementing it are still unclear (65).

Consuming a healthy diet, such as the Mediterranean diet (66-70), a prudent diet (71), the Dietary Approaches to Stop Hypertension (DASH) diet (69, 70) and the Mediterranean-DASH Intervention for Neurodegenerative Delay (MIND) diet (70), can also improve cognitive performance. The Mediterranean diet and the prudent diet are characterized by high fruit, vegetable, fish, wholegrain, oil, legume, nut and low-fat dairy intake while limiting the intake of red meat, sweets, sugar-containing beverages, fat and cholesterol (69, 71). The DASH diet is largely similar but places a larger emphasis on low-fat dairy and low dietary sodium intake and alcohol abstinence (69), whereas the MIND diet combines the Mediterranean and DASH diets and emphasizes plant-based foods over animal foods and foods high in saturated fat (70).

### *Nootropics*

Nootropics are pharmacological substances that can be used to enhance cognitive performance, such as (ar)modafinil and methylphenidate, but also caffeine or nicotine (72-74). They are popular in the general population (mainly among students in order to enhance their academic performance). Taking nootropics is a much easier way to enhance cognitive performance compared to strategies such as exercise or consuming a healthy diet, but it is associated with considerable drawbacks. Firstly, substances such as methylphenidate or modafinil (primarily indicated in The Netherlands for attention deficit / hyperactivity disorder and for narcolepsy, respectively (75, 76)) were not developed with the aim of cognitive

enhancement in mind. As such, the evidence base for their cognitive performance-enhancing properties is weak, especially in people with comparable demographical and pathophysiological characteristics as typical patients with COPD. Most meta-analyses and systematic reviews have investigated their effects in younger adults or in patients with traumatic brain injury (77, 78). Secondly, nootropics can have unintended and undesirable side effects such as headaches or insomnia (75, 76). Thirdly, they should be very carefully administered as overdosing can lead to suboptimal cognitive performance (79) and harmful effects on cerebral functioning (80). Therefore, using nootropics with the sole aim of cognitive enhancement (i.e., without any other indication) cannot be recommended.

One class of medication that might be feasible as a nootropic in patients with COPD are phosphodiesterase (PDE)-4 inhibitors such as roflumilast. They have been shown to improve cognitive performance in animal models of Alzheimer's disease (81) and chronic cerebral hypoperfusion (82). Acute administration of 100µg (but not 250, 300, 500 or 1000µg) roflumilast significantly improved immediate recall memory performance in healthy young adults (83) and delayed recall memory performance in healthy older adults (84).

Roflumilast has recently been indicated in The Netherlands to treat patients with severe COPD ( $FEV_1 < 50\%$  predicted) who suffer from frequent exacerbations despite optimal triple therapy (85). Its potential to simultaneously improve the health status and cognitive functioning of patients with COPD would make it a promising intervention. Moreover, it is relatively cheap, with an estimated price per day of €1.74 (85). However, although acute administration of the recommended therapeutic dose for COPD (500µg) tended to lead to better cognitive performance compared to placebo in participants without COPD, these differences were not significant (83, 84). The effects of long-term roflumilast administration on human cognitive performance have not been investigated to our knowledge. Hence, whether roflumilast administration alone or as part of standard COPD management could contribute to cognitive improvement remains to be investigated.

### *Pulmonary rehabilitation*

PR is also prescribed to patients with severe COPD as part of standard disease management (86). PR includes many components with proven benefits on cognitive performance, such as exercise training and nutritional support (86), and the available evidence indeed provides tentative indications of favorable effects of PR on cognitive performance (43, 87-89). The necessary investment in terms of money, time and effort do not make PR a viable option to improve cognitive performance of patients with COPD on a large scale, but it could benefit those

patients who are referred for PR based on the existing indications. Moreover, the likely poor cost-effectiveness of PR to improve cognitive functioning does not negate the fact that incorporating its key elements, such as physical activity, exercise and dietary improvement, into daily life will likely benefit cognitive performance, as discussed above. Future research should expand on the existing evidence base with methodologically stronger studies. They should for instance, in contrast to earlier studies, include a non-PR control group (43, 87) and include a broader sample of patients than only certain subsets such as those who have undergone a lung transplantation (89).

PR is a very effective intervention to improve health outcomes, but its transfer into an actual healthier lifestyle after completion of the program and after the associated support and structure are taken away, is only poor (90). Interventions targeting patients' daily lives in the first place may be more effective.

### *Comprehensive lifestyle interventions*

Many modifiable lifestyle factors influence cognitive performance. The Dutch Alzheimer Centre lists mental and physical inactivity, smoking, depression, overweight, hypertension and diabetes as important risk factors to develop dementia (91). Targeting one factor in an intervention while leaving all others constant therefore likely only has a small effect on cognitive performance. Instead, interventions likely become more effective as more factors are targeted.

However, in spite of this enhanced effectiveness in theory, the current empirical evidence for such multidomain interventions is mixed. Over the past decade, three large-scale studies have investigated the potential of comprehensive lifestyle interventions to diminish the risk of developing dementia: the Finnish Geriatric Intervention Study to Prevent Cognitive Impairment and Disability (FINGER) (92), the preDIVA study (93) and the Multidomain Alzheimer Preventive Trial (MAPT) (94). The 2-year FINGER strategy consisted of nutritional management, exercise training, cognitive training and vascular risk monitoring. The 6-year preDIVA study combined usual cardiovascular care with tailored lifestyle advice based on motivational interviewing principles; the 3-year MAPT trial offered cognitive training, physical activity, nutritional supplementation and educational sessions. The FINGER trial was able to prevent cognitive decline to a significantly greater degree than general health advice, but the preDIVA and MAPT studies did not lead to significant changes in cognitive decline compared to placebo groups. In conclusion, there is a clear need for future studies to investigate the potential of such multidomain interventions, and to identify which components are effective to which degree.

## Working memory training and adherence to a healthy lifestyle

In this thesis WMT was hypothesized to be able to contribute to a healthier lifestyle through its ability to increase cognitive control (95, 96). This would enable patients to inhibit automatic responses in favor of more opportune controlled ones (97). However, the Cogtrain trial showed no effects of WMT on adherence to a healthy lifestyle, the recall of prespecified healthy lifestyle goals or motivation towards a healthy lifestyle.

One potential explanation for these non-significant results is that transfer from improved performance on one cognitive domain to other cognitive domains is already scarce (31, 32). Transfer towards healthy lifestyle outcomes would have to reach even further. All of these outcomes are determined by a potentially endless number of pathways involving motivational, environmental (e.g., the weather, social support, financial issues) and health-related factors (e.g., disease progression). WMT-induced improved cognitive control may contribute to adherence to a healthy lifestyle, but it is not a panacea.

Secondly, the positive effects of earlier studies investigating the effects of WMT on healthy lifestyle outcomes or stress susceptibility were largely found on lab-based parameters or tasks (such as a bogus taste test in participants with overweight or metabolic parameters in those with type 2 diabetes) instead of actual healthier lifestyles (i.e., weight loss or disease management) (95, 98). Even studies which selectively included specific subgroups of participants who stood to gain a lot by participating, such as people with overweight in studies with weight loss as a primary outcome (99) or heavy drinkers in a study aiming to reduce substance abuse (100), were largely ineffective. The one recent study investigating the effects of WMT on stress responses found positive results on the biological response to stress (measured by cortisol levels) but did not include a measure of subjective stress perception (101).

Thirdly, increasing cognitive control might not be necessary or sufficient to establish a healthy lifestyle. It might not be sufficient because a large proportion of patients with COPD appear not to know what is and is not healthy: 30% of them qualify as being health illiterate (102, 103). Future studies and clinical efforts may therefore have to add educational components in order to enable patients to keep the right information in mind. However, the effectiveness of combining only WMT and education is doubtful (95), implying that these components will also have to be part of a more comprehensive interventional strategy.

Combining education, WMT and additional interventional components is a highly cumbersome way to achieve a healthier lifestyle. All of this might not be necessary in the first place, because it could simply and effectively be bypassed using a technique called 'nudging' (104). Nudging consists of altering certain aspects of people's environment with the aim of automatically guiding them towards a certain choice even though they retain the freedom to choose any number of other options (104). It therefore minimizes the necessary involvement of cognitive control, instead of maximizing its capacity. Nudging can be very easy and subtle, and it can unconsciously change behavior: it can consist of placing the fruit and vegetables department immediately at the entrance of a supermarket and making it impossible not to pass through it before reaching other departments with unhealthier items. Importantly, customers retain full freedom to pick unhealthy products, but the likelihood of them buying fruit and vegetables is much larger compared to when the fruit and vegetables department is located in the middle or the back of the store. Its non-coercive nature also makes nudging much more likely to be effective, also in the longer term, compared to educational or prescriptive approaches telling you what you 'should' or 'should not' do (105, 106).

## General conclusion

CI is an important comorbidity of COPD, which should be better recognized and dealt with in the clinic. WMT as single intervention does not improve cognitive performance in COPD. Future initiatives to improve cognitive performance in patients with COPD should focus on alternative, preferably multimodal, interventions.

## References

1. Sievi N, Brack T, Brutsche MH, Frey M, Irani S, Leuppi JD, et al. Physical activity declines in COPD while exercise capacity remains stable: a longitudinal study over 5 years. *Respiratory Medicine*. 2018;141:1-6.
2. Clarenbach CF, Sievi N, Haile SR, Brack T, Brutsche MH, Frey M, et al. Determinants of annual change in physical activity in COPD. *Respirology*. 2017;22(6):1133-9.
3. Laudisio A, Costanzo L, Di Gioia C, Delussu AS, Traballese M, Gemma A, et al. Dietary intake of elderly outpatients with chronic obstructive pulmonary disease. *Archives of Gerontology and Geriatrics*. 2016;64:75-81.
4. Chambaneau A, Filaire M, Jubert L, Bremond M, Filaire E. Nutritional intake, physical activity and quality of life in COPD patients. *International Journal of Sports Medicine*. 2016;37:730-7.
5. Van de Bool C, Mattijssen-Verdonschot C, Van Melick PPMJ, Spruit MA, Franssen FME, Wouters EFM, et al. Quality of dietary intake in relation to body composition in patients with chronic obstructive pulmonary disease eligible for pulmonary rehabilitation. *European Journal of Clinical Nutrition*. 2014;68(2):159-65.
6. Yohannes AM, Chen W, Moga AM, Leroi I, Connolly MJ. Cognitive impairment in chronic obstructive pulmonary disease and chronic heart failure: a systematic review and meta-analysis of observational studies. *JAMDA*. 2017;18:451.e1-e11.
7. Higbee DH, Dodd JW. Cognitive impairment in COPD: an often overlooked comorbidity. *Expert Review of Respiratory Medicine*. 2020;in press.
8. Dal Negro RW, Bonadiman L, Tognella S, Bricolo FP, Turco P. Extent and prevalence of cognitive dysfunction in chronic obstructive pulmonary disease, chronic non-obstructive bronchitis, and in asymptomatic smokers, compared to normal reference values. *International Journal of COPD*. 2014;26(9):675-83.
9. Antonelli Incalzi R, Gemma A, Marra C, Capparella O, Fuso L, Carbonin P. Verbal memory impairment in COPD: its mechanisms and clinical relevance. *Chest*. 1997;112(6):1506-13.
10. Dodd JW, Getov SV, Jones PW. Cognitive function in COPD. *European Respiratory Journal*. 2010;35:913-22.
11. Chang SS, Chen S, McAvay GJ, Tinetti ME. Effect of coexisting chronic obstructive pulmonary disease and cognitive impairment on health outcomes in older adults. *Journal of the American Geriatric Society*. 2012;60(10):1839-46.
12. Disler RT, Spiliopoulos N, Inglis SC, Currow DC, Davidson PM. Cognitive screening in chronic obstructive pulmonary disease: patient's perspectives. *Disability and Rehabilitation*. 2020;42(9):1233-9.
13. Global Initiative for Chronic Obstructive Lung Disease (GOLD). Global Strategy for the Diagnosis, Management and Prevention of Chronic Obstructive Pulmonary Disease. Agustí A, Vogelmeier C, editors 2020.



14. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": a practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*. 1975;12(3):189-98.
15. Villeneuve S, Pepin V, Rahayel S, Bertrand J-A, De Lorimier M, Rizk A, *et al*. Mild cognitive impairment in moderate to severe COPD. *Chest*. 2012;142(6):1516-23.
16. Tsoi KKF, Chan JYC, Hirai HW, Wong SYS, Kwok TCY. Cognitive tests to detect dementia: a systematic review and meta-analysis. *JAMA Internal Medicine*. 2015;175(9):1450-8.
17. Benz E, Trajanoska K, Lahousse L, Schoufour JD, Terzikhan N, De Roos E, *et al*. Sarcopenia in COPD: a systematic review and meta-analysis. *European Respiratory Review*. 2019;28:190049.
18. Cebron Lipovec N, Beijers RJ, Van den Borst B, Doehner W, Lainscak M, Schols AMWJ. The prevalence of metabolic syndrome in chronic obstructive pulmonary disease: a systematic review. *COPD*. 2016;13(3):399-406.
19. Cavalieri M, Ropele S, Petrovic K, Pluta-Fuerst A, Homayoon N, Enzinger C, *et al*. Metabolic syndrome, brain magnetic resonance imaging, and cognition. *Diabetes Care*. 2010;33(12):2489-95.
20. Falkowski J, Atchison T, DeButte-Smith M, Weiner MF, O'Bryant S. Executive functioning and the metabolic syndrome: a Project FRONTIER study. *Archives of Clinical Neuropsychology*. 2014;29:47-53.
21. Peng T-C, Chen W-L, Wu L-W, Chang Y-W, Kao T-W. Sarcopenia and cognitive impairment: a systematic review and meta-analysis. *Clinical Nutrition*. 2019;S0261-5614(19):33200-5.
22. Cabett Cipolli G, Sandes Yassuda M, Aprahamian I. Sarcopenia is associated with cognitive impairment in older adults: a systematic review and meta-analysis. *Journal of Nutrition, Health and Aging*. 2019;23:525-31.
23. Moon JH, Moon JH, Kim KM, Choi SH, Lim S, Park KS, *et al*. Sarcopenia as a predictor of future cognitive impairment in older adults. *Journal of Nutrition, Health and Aging*. 2016;20(5):496-502.
24. Mercado N, Ito K, Barnes PJ. Accelerated ageing of the lung in COPD: new concepts. *Thorax*. 2015;70:482-9.
25. Martin-Ruiz C, Jagger C, Kingston A, Collerton J, Catt M, Davies K, *et al*. Assessment of a large panel of candidate biomarkers of ageing in the Newcastle 85+ study. *Mechanisms of Ageing and Development*. 2011;132(10):496-502.
26. Yaffe K, Lindquist K, Kluse M, Cawthon R, Harris T, Hsueh W-C, *et al*. Telomere length and cognitive function in community-dwelling elders: findings from the Health ABC study. *Neurobiology of Aging*. 2011;32(11):2055-60.
27. Hägg S, Zhan Y, Karlsson R, Gerritsen L, Ploner A, Van der Lee SJ, *et al*. Short telomere length is associated with impaired cognitive performance in European ancestry cohorts. *Translational Psychiatry*. 2017;7:e1100.

28. Devore EE, Prescott J, De Vivo I, Grodstein F. Relative telomere length and cognitive decline in the Nurses' Health Study. *Neuroscience Letters*. 2011;492(1):15-8.
29. Rask L, Bendix L, Harbo M, Fagerlund B, Mortensen EL, Lauritzen MJ, et al. Cognitive change during the life course and leukocyte telomere length in late middle-aged men. *Frontiers in Aging Neuroscience*. 2016;8:300.
30. Conway ARA, Kane MJ, Engle RW. Working memory capacity and its relation to general intelligence. *Trends in Cognitive Sciences*. 2003;7(12):547-52.
31. Melby-Lervag M, Hulme C. Is working memory training effective? A meta-analytic review. *Developmental Psychology*. 2013;49(2):270-91.
32. Melby-Lervag M, Redick TS, Hulme C. Working memory training does not improve performance on measures of intelligence or other measures of "far transfer": evidence from a meta-analytic review. *Perspectives on Psychological Science*. 2016;11(4):512-34.
33. Teixeira-Santos AC, Moreira CS, Magalhaes R, Magalhaes C, Pereira DR, Leite J, et al. Reviewing working memory training gains in healthy older adults: a meta-analytic review of transfer for cognitive outcomes. *Neuroscience and Biobehavioral Reviews*. 2019;103:163-77.
34. Antonelli Incalzi R, Corsonello A, Trojano L, Pedone C, Acanfora D, Spada A, et al. Cognitive training is ineffective in hypoxemic COPD: a six-month randomized controlled trial. *Rejuvenation Research*. 2008;11:239-50.
35. Souders DJ, Boot WR, Blocker K, Vitale T, Roque NA, Charness N. Evidence for narrow transfer after short-term cognitive training in older adults. *Frontiers in Aging Neuroscience*. 2017;9:41.
36. Lillo-Crespo M, Forner-Ruiz M, Riquelme-Galindo J, Ruiz-Fernández D, García-Sanjuan S. Chess practice as a protective factor in dementia. *International Journal of Environmental Research and Public Health*. 2019;16(12):2116.
37. Ferreira N, Owen A, Mohan A, Corbett A, Ballard C. Associations between cognitively stimulating leisure activities, cognitive function and age-related cognitive decline. *International Journal of Geriatric Psychiatry*. 2015;30(4):422-30.
38. Kelly ME, Duff H, Kelly S, McHugh Power JE, Brennan S, Lawlor BA, et al. The impact of social activities, social networks, social support and social relationships on the cognitive functioning of healthy older adults: a systematic review. *Systematic Reviews*. 2017;6:259.
39. Biddle KD, D'Oleire Uquillas F, Jacobs HIL, Zide B, Kirn DR, Rentz DM, et al. Social engagement and amyloid- $\beta$ -related cognitive decline in cognitively normal older adults. *American Journal of Geriatric Psychiatry*. 2019;27(11):1247-56.
40. Fernandes RM, Correa MG, Dos Santos MAR, Almeida APCPSC, Fagundes NCF, Maia LC, et al. The effects of moderate physical exercise on adult cognition: a systematic review. *Frontiers in Physiology*. 2018;9:667.

41. Falck RS, Davis JC, Best JR, Crockett RA, Liu-Ambrose T. Impact of exercise training on physical and cognitive function among older adults: a systematic review and meta-analysis. *Neurobiology of Aging*. 2019;79:119-30.
42. Song D, Yu DSF, Li PWC, Lei Y. The effectiveness of physical exercise on cognitive and psychological outcomes in individuals with mild cognitive impairment: a systematic review and meta-analysis. *International Journal of Nursing Studies*. 2018;79:155-64.
43. Bonnevie T, Medrinal C, Combret Y, Debeaumont D, Lamia B, Muir J-F, *et al.* Mid-term effects of pulmonary rehabilitation on cognitive function in people with severe chronic obstructive pulmonary disease. *International Journal of Chronic Obstructive Pulmonary Disease*. 2020;15:1111-21.
44. Pedersen BK. Physical activity and muscle-brain crosstalk. *Nature Reviews Endocrinology*. 2019;15:383-92.
45. Chen HT, Chung YC, Chen YJ, Ho SY, Wu HJ. Effects of different types of exercise on body composition, muscle strength, and IGF-1 in the elderly with sarcopenic obesity. *Journal of the American Geriatric Society*. 2017;65(4):827-32.
46. Cho S-Y, Roh H-T. Taekwondo enhances cognitive function as a result of increased neurotrophic growth factors in elderly women. *International Journal of Environmental Research and Public Health*. 2019;16(6):962.
47. Woost L, Bazin P-L, Taubert M, Trampel R, Tardif CL, Garthe A, *et al.* Physical exercise and spatial training: a longitudinal study of effects on cognition, growth factors, and hippocampal plasticity. *Nature Scientific Reports*. 2018;8:4239.
48. Bettio L, Thacker JS, Hutton C, Christie BR. Modulation of synaptic plasticity by exercise. *International Review of Neurobiology*. 2019;147:295-322.
49. Joisten N, Kummerhoff F, Koliamitra C, Schenk A, Walzik D, Hardt L, *et al.* Exercise and the kynurenine pathway: current state of knowledge and results from a randomized cross-over study comparing acute effects of endurance and resistance training. *Exercise Immunology Review*. 2020;26:24-42.
50. Valente-Silva P, Ruas JL. Tryptophan-kynurenine metabolites in exercise and mental health. In: Spiegelman B, editor. *Hormones, Metabolism and the Benefits of Exercise. Research and Perspectives in Endocrine Interactions*: Springer; 2017.
51. Joisten N, Walzik D, Metcalfe AJ, Bloch W, Zimmer P. Physical exercise as kynurenine pathway modulator in chronic diseases: implications for immune and energy homeostasis. *International Journal of Tryptophan Research*. 2020;13:1178646920938688.
52. Hafstad Solvang S-E, Nordrehaug JE, Tell GS, Nygård O, McCann A, Ueland PM, *et al.* The kynurenine pathway and cognitive performance in community-dwelling older adults: the Hordaland Health Study. *Brain, Behavior and Immunity*. 2019;755:155-62.
53. Agudelo LZ, Femenía T, Orhan F, Porsmyr-Palmertz M, Gojny M, Martínez-Redondo V, *et al.* Skeletal muscle PGC-1 $\alpha$ 1 modulates kynurenine metabolism and mediates resilience to stress-induced depression. *Cell*. 2014;159(1):33-45.

54. Gosker HR, Clarke G, De Theije CC, Cryan JF, Schols AM. Impaired skeletal muscle kynurenine metabolism in patients with chronic obstructive pulmonary disease. *Journal of Clinical Medicine*. 2019;8(7):915.
55. Dinan TG, Cryan JF. The microbiome-gut-brain axis in health and disease. *Gastroenterology Clinics of North America*. 2017;46(1):77-89.
56. Strandwitz P. Neurotransmitter modulation by the gut microbiota. *Brain Research*. 2018;1693:128-33.
57. Sun L, Paräkylä J, Holm K, Haapasalo J, Lehtimäki K, Ogawa KH, et al. Vagus nerve stimulation improves working memory performance. *Journal of Clinical and Experimental Neuropsychology*. 2017;39(10):954-64.
58. Colzato L, Beste C. A literature review on the neurophysiological underpinnings and cognitive effects of transcutaneous vagus nerve stimulation: challenges and future directions. *Journal of Neurophysiology*. 2020;123(5):1739-55.
59. Long-Smith C, O'Riordan KJ, Clarke G, Stanton C, Dinan TG, Cryan JF. Microbiota-gut-brain axis: new therapeutic opportunities. *Annual Review of Pharmacology and Toxicology*. 2020;60:477-502.
60. Calder PC. Omega-3 fatty acids and inflammatory processes: from molecules to man. *Biochemical Society Transactions*. 2017;45(5):1105-15.
61. Schwarcz R, Stone TW. The kynurenine pathway and the brain: challenges, controversies and promises. *Neuropharmacology*. 2017;112:237-47.
62. García-Esquinas E, Ortolá R, Banegas JR, Lopez-García E, Rodríguez-Artalejo F. Dietary n-3 polyunsaturated fatty acids, fish intake and healthy ageing. *International Journal of Epidemiology*. 2019;48(6):1914-24.
63. Bo Y, Zhang X, Wang Y, You J, Cui H, Zhu Y, et al. The n-3 polyunsaturated fatty acids supplementation improved the cognitive function in the Chinese elderly with mild cognitive impairment: a double-blind randomized controlled trial. *Nutrients*. 2017;9(1):54.
64. Berding K, Long-Smith C, Carbia C, Bastiaansen TFS, Van de Wouw M, Wiley N, et al. A specific dietary fibre supplementation improves cognitive performance - an exploratory randomised, placebo-controlled, crossover study. *Psychopharmacology*. 2020;In press.
65. Goodwill AM, Szoëke C. A systematic review and meta-analysis of the effect of low vitamin D on cognition. *Journal of the American Geriatric Society*. 2017;65(10):2161-8.
66. Martínez-Lapiscina EH, Clavero P, Toledo E, Estruch R, Salas-Salvadó J, San Julián B, et al. Mediterranean diet improves cognition: the PREDIMED-NAVARRA randomised trial. *Journal of Neurology, Neurosurgery and Psychiatry*. 2013;84(12):1318-25.
67. Petersson SD, Philippou E. Mediterranean diet, cognitive function, and dementia: a systematic review of the evidence. *Advances in Nutrition*. 2016;7(5):889-904.

68. Aridi YS, Walker JL, Wright ORL. The association between the Mediterranean dietary pattern and cognitive health: a systematic review. *Nutrients*. 2017;9(7):674.
69. McGrattan AM, McGuinness B, McKinley MC, Kee F, Passmore P, Woodside JV, *et al.* Diet and inflammation in cognitive ageing and Alzheimer's disease. *Current Nutrition Reports*. 2019;8(2):53-65.
70. Van den Brink AC, Brouwer-Brolsma EM, Berendsen AAM, Van de Rest O. The Mediterranean, Dietary Approaches to Stop Hypertension (DASH), and Mediterranean-DASH Intervention for Neurodegenerative Delay (MIND) diets are associated with less cognitive decline and a lower risk of Alzheimer's disease - a review. *Advances in Nutrition*. 2019;10(6):1040-65.
71. Shakersain B, Santoni G, Larsson SC, Faxen-Irving G, Fastborn J, Fratiglioni L, *et al.* Prudent diet may attenuate the adverse effects of Western diet on cognitive decline. *Alzheimer's & Dementia*. 2016;12(2):100-9.
72. Battleday RM, Brem A-K. Modafinil for cognitive neuroenhancement in healthy non-sleep-deprived subjects: a systematic review. *European Neuropsychopharmacology*. 2015;25:1865-81.
73. Ilieva IP, Hook CJ, Farah MJ. Prescription stimulants' effects on healthy inhibitory control, working memory, and episodic memory: a meta-analysis. *Journal of Cognitive Neuroscience*. 2015;27(6):1069-89.
74. Heishman SJ, Kleykamp BA, Singleton EG. Meta-analysis of the acute effects of nicotine and smoking on human performance. *Psychopharmacology*. 2010;210:453-69.
75. Methylfenidaat: Zorginstituut Nederland; [cited 2020 October 5]. Available from: <https://www.farmacotherapeutischkompas.nl/bladeren/preparaatteksten/m/methylfenidaat>.
76. Modafinil: Zorginstituut Nederland; [cited 2020 October 5]. Available from: <https://www.farmacotherapeutischkompas.nl/bladeren/preparaatteksten/m/modafinil>.
77. Huang C-H, Huang C-C, Sun C-K, Lin G-H, Hou W-H. Methylphenidate on cognitive improvement in patients with traumatic brain injury: a meta-analysis. *Current Neuropharmacology*. 2016;14(3):272-81.
78. Barnett M, Reid L. The effectiveness of methylphenidate in improving cognition after brain injury in adults: a systematic review. *Brain Injury*. 2020;34(1):1-10.
79. Wood S, Sage JR, Anagnostaras SG. Psychostimulants and cognition: a continuum of behavioral and cognitive activation. *Pharmacological Reviews*. 2014;66(1):193-221.
80. Urban KR, Gao W-J. Performance enhancement at the cost of potential brain plasticity: neural ramifications of nootropic drugs in the healthy developing brain. *Frontiers in Systems Neuroscience*. 2014;8:38.
81. Feng H, Wang C, He W, Wu X, Li S, Zeng Z, *et al.* Roflumilast ameliorates cognitive impairment in APP/PS1 mice via cAMP/CREB/BDNF signaling and anti-neuroinflammatory effects. *Metabolic Brain disease*. 2019;34:583-91.

82. Santiago A, Mendes Soares L, Schepers M, Milani H, Vanmierlo T, Prickaerts J, *et al.* Roflumilast promotes memory recovery and attenuates white matter injury in aged rats subjected to chronic cerebral hypoperfusion. *Neuropharmacology*. 2018;138:360-70.
83. Van Duinen MA, Sambeth A, Heckman PRA, Smit S, Tsai M, Lahu G, *et al.* Acute administration of roflumilast enhances immediate recall of verbal word memory in healthy young adults. *Neuropharmacology*. 2018;131:31-8.
84. Blokland A, Van Duinen MA, Sambeth A, Heckman PRA, Tsai M, Lahu G, *et al.* Acute treatment with the PDE4 inhibitor roflumilast improves verbal word memory in healthy old individuals: a double-blind placebo-controlled study. *Neurobiology of Aging*. 2019;77:37-43.
85. Roflumilast: Zorginstituut Nederland; [cited 2020 September 28]. Available from: <https://www.farmacotherapeutischkompas.nl/bladeren/preparaatteksten/r/roflumilast>.
86. Spruit MA, Singh SJ, Garvey C, ZuWallack R, Nici L, Rochester C, *et al.* An official American Thoracic Society / European Respiratory Society statement: Key concepts and advances in pulmonary rehabilitation. *American Journal of Respiratory and Critical Care Medicine*. 2013;188(8):e13-e64.
87. Emery CF, Leatherman NE, Burkner EJ, MacIntyre NR. Psychological outcomes of a pulmonary rehabilitation program. *Chest*. 1991;100(3):613-7.
88. Pereira EDB, Sampaio Viana C, Taunay TCE, Sales PU, Lima JWO, Holanda MA. Improvement of cognitive function after a three-month pulmonary rehabilitation program for COPD patients. *Lung*. 2011;189:279-85.
89. Andrianopoulos V, Gloeckl R, Boensch M, Hoster K, Schneeberger T, Jarosch I, *et al.* Improvements in functional and cognitive status following short-term pulmonary rehabilitation in COPD lung transplant patients: a pilot study. *ERJ Open Research*. 2019;5(3):00060-2019.
90. Troosters T, Demeyer H, Hornikx M, Camillo CA, Janssens W. Pulmonary rehabilitation. *Clinics in Chest Medicine*. 2014;35(1):241-9.
91. Het voorkomen van dementie: Alzheimer Centrum Nederland; n.d. [Available from: <https://www.alzheimer-nederland.nl/dementie/oorzaken-preventie/voorkomen>].
92. Ngandu T, Lehtisalo J, Solomon A, Levälähti E, Ahtiluoto S, Antikainen R, *et al.* A 2 year multidomain intervention of diet, exercise, cognitive training, and vascular risk monitoring versus control to prevent cognitive decline in at-risk elderly people (FINGER): a randomised controlled trial. *Lancet*. 2015;385(9984):2255-63.
93. Moll van Charante EP, Richard E, Eurelings LS, Van Dalen J-W, Ligthart SA, Van Bussel EF, *et al.* Effectiveness of a 6-year multidomain vascular care intervention to prevent dementia (preDIVA): a cluster-randomised controlled trial. *Lancet*. 2016;388(10046):797-805.

94. Andrieu S, Guyonnet S, Coley N, Cantet C, Bonnefoy M, Bordes S, *et al.* Effect of long-term omega 3 polyunsaturated fatty acid supplementation with or without multidomain intervention on cognitive function in elderly adults iwth memory complaints (MAPT): a randomised, placebo-controlled trial. *Lancet Neurology*. 2017;16(5):377-89.
95. Dassen FCM, Houben K, Van Breukelen GJP, Jansen A. Gamified working memory training in overweight individuals reduces food intake but not body weight. *Appetite*. 2018;124:89-98.
96. Kahneman D. A perspective on judgment and choice: mapping bounded rationality. *American Psychologist*. 2003;58(9):697-720.
97. Diamond A. Executive functions. *Annual Review of Psychology*. 2013;64:135-68.
98. Whitelock V, Nouwen A, Houben K, Van den Akker O, Rosenthal M, Higgs S. Does working memory training improve dietary self-care in type 2 diabetes mellitus? Results of a double blind randomised controlled trial. *Diabetes Research and Clinical Practice*. 2018;143:204-14.
99. Houben K, Dassen FCM, Jansen A. Taking control: working memory training in overweight individuals increasesself-regulation of food intake. *Appetite*. 2016;105:567-74.
100. Houben K, Wiers RW, Jansen A. Getting a grip on drinking behavior: training working memory to reduce alcohol abuse. *Psychological Science*. 2011;22(7):968-75.
101. Jopling E, Gotlib IH, LeMoult J. Effects of working memory training on cognitive, affective and biological responses to stress in major depression: a novel cognitive bias modification protocol. *Journal of Affective Disorders*. 2020;265:45-51.
102. O'Connor R, Muellers KA, Arvanitis M, Vicencio DP, Wolf MS, Wisnivesky JP, *et al.* Effects of health literacy and cognitive abilities on COPD self-management behaviors: a prospective cohort study. *Respiratory Medicine*. 2019;160:105630.
103. Muellers KA, Chen L, O'Connor R, Wolf MS, Federman AD, Wisnivesky JP. Health literacy and medication adherence in COPD patients: when caregiver presence is not sufficient. *COPD*. 2019;16(5-6):362-7.
104. Thaler RH, Sunstein CR. *Nudge: improving decisions about health, wealth and happiness*. New Haven, CT: Yale University Press; 2008.
105. Deci EL, Ryan RM. *Intrinsic motivation and self-determination in human behavior*. New York, NY: Plenum; 1985.
106. Markland D, Ryan RM, Tobin VJ, Rollnick S. Motivational interviewing and self-determination theory. *Journal of Social and Clinical Psychology*. 2005;24(6):811-31.





# Summary

Chronic obstructive pulmonary disease (COPD) is a high-burden chronic respiratory disease. It is currently the third most common cause of death worldwide, with an estimated 3 million deaths in 2016. Traditional interventions to improve the health status of patients with COPD, such as supervised pulmonary rehabilitation (PR) programs, are effective in the short term but often do not translate into a healthier and generally more active lifestyle after completion of the program. Promising additional or alternative approaches to maintain or improve health status of COPD patients include motivational counseling towards a healthy lifestyle, targeted nutritional intervention or working memory training (WMT).

Cognitive impairment (CI) is a common extrapulmonary manifestation of COPD, and is evidenced for example by memory loss or reduced speed of thinking. CI is a problem in itself, but also affects overall health status of patients with COPD and may influence their ability to adhere to a healthy lifestyle.

The main aim of this thesis was therefore to investigate determinants of cognitive performance in patients with COPD and the potential efficacy of WMT next to motivational regulations in increasing patients' adherence to a healthy lifestyle.

**Chapter 2** reports the long-term results of the randomized, placebo-controlled NUTRAIN trial. This study investigated the cost-effectiveness of a nutritional management strategy in patients with COPD and low muscle mass (often referred to as sarcopenia). The strategy consisted of targeted nutritional supplementation (leucine, polyunsaturated fatty acids and vitamin D) in a medical drink and dietary counselling during a 4-month outpatient PR program and a subsequent 8-month maintenance phase. Relative to usual care (PR without follow-up intervention) physical capacity increased in both groups, but the nutritional management approach ameliorated plasma levels of the supplemented nutrients, prevented unintended weight loss, and positively influenced daily physical activity and generic health.

**Chapter 3** investigates whether motivational regulations are related to healthy lifestyle behaviors (physical activity and adherence to a healthy diet) and the responsiveness thereof after an outpatient PR program. Levels of identified motivation (i.e., performing an action because it is valued as personally important to someone) predicted dietary quality change during PR and its levels after PR; levels of introjected motivation (i.e., performing an action out of fear of the shame associated with not doing it) predicted change in physical activity levels during PR and its levels after PR.

**Chapter 4** provides a narrative review of the initial knowledge base around CI in COPD. Based on the current literature, several disease-specific factors, comorbidities and unhealthy lifestyle factors may contribute to the emergence of CI in COPD, which affects around one in three patients. A diffuse pattern of cognitive domains is affected, in contrast to disease-specific patterns such as the selective loss of memory (but not other cognitive domains) in Alzheimer's disease. These impairments have a detrimental impact on patients' self-management abilities, health outcomes, ability to engage in activities of daily living and quality of life, while increasing their hospitalization risk.

**Chapter 5** investigates whether the presence of the metabolic syndrome or sarcopenia is related to an elevated prevalence and/or different pattern of CI in patients with COPD referred for PR. These conditions are more common in COPD than in healthy controls and adversely affect health status and survival. COPD patients with one of these comorbidities might be especially vulnerable to develop CI, as these conditions have been independently associated with poor cognitive functioning in non-COPD populations. In the studied COPD population referred for PR, the prevalence of metabolic syndrome (54.7%) and sarcopenia (30.0%) was high, but not associated with a higher prevalence of CI.

**Chapter 6** reports the results of the randomized, placebo-controlled Cogtrain trial. In this trial a 12-week home-based working memory training (WMT) program was administered, followed by a 12-week maintenance program. The efficacy of WMT to establish and maintain improvements in cognitive performance, adherence to a healthy lifestyle and cognitive stress susceptibility was investigated in patients with COPD. Although the intervention improved performance on the actual (working memory) tasks that were used for training, these improvements did not generalize to other cognitive performance domains. The WMT did not improve adherence to a healthy lifestyle or cognitive stress susceptibility either. These results indicate that WMT is not an effective way to improve cognitive performance and potentially related healthy lifestyle behavior in patients with COPD.

**Chapter 7** positions the results of the current thesis in a broader scientific and clinical perspective, with particular emphasis on cognitive functioning in COPD and on the potential efficacy of interventions to improve cognitive performance and adherence to a healthy lifestyle in patients with COPD.

The overall results of the current thesis indicate that CI is an important comorbidity of COPD, which should be better recognized and dealt with in clinical care. CI has an independent substantial detrimental impact on daily-life functioning and health outcomes of patients with COPD. Metabolic abnormalities (i.e., the

## Summary

metabolic syndrome and sarcopenia) do not appear to be useful screening factors for CI in patients with COPD. WMT did not improve cognitive functioning or adherence to a healthy lifestyle. However, motivational regulations were shown to be potentially effective interventional targets to establish and maintain a healthy lifestyle following PR.





# Samenvatting

COPD is een veelvoorkomende en ernstige chronische longziekte, die momenteel de op twee na meest voorkomende doodsoorzaak wereldwijd is. In 2016 was COPD naar schatting verantwoordelijk voor 3 miljoen doden. Gangbare niet-medicamenteuze interventies om de algemene gezondheid van patiënten met COPD te verbeteren, zoals longrevalidatie, zijn effectief op korte termijn, maar beklijven vaak niet op de langere termijn via een gezondere leefstijl. Een gerichte voedingsaanpak, motivationele counseling en werkgeheugentraining zijn veelbelovende aanvullende of alternatieve behandelingen om de gezondheid en leefstijl van patiënten met COPD te verbeteren.

Verminderd cognitief functioneren, zoals bijvoorbeeld geheugenverlies of een vertraagd denkvermogen, komt vaak voor bij COPD. Deze cognitieve beperkingen beïnvloeden de algehele gezondheid van patiënten met COPD en verminderen hun vermogen om gezond te leven.

In deze thesis is nader onderzoek verricht naar de risicofactoren voor cognitief functieverlies bij COPD alsmede de trainbaarheid van het cognitief functioneren middels werkgeheugentraining. Daarnaast is onderzocht of deze training naast een doelgerichte voedingsinterventie en inzicht in verschillende vormen van motivatie kan bijdragen aan een gezondere leefstijl van patiënten met COPD.

**Hoofdstuk 2** rapporteert de langetermijnsresultaten van de gerandomiseerde, placebo-gecontroleerde NUTRAIN-trial. De interventiegroep ontving een medische drinkvoeding die verrijkt was met leucine, meervoudig onverzadigde vetzuren en vitamine D gedurende een 4 maanden durend poliklinisch revalidatieprogramma en counseling middels motivationele interviews tijdens de daaropvolgende onderhoudsfase van 8 maanden. Deze interventie verbeterde na 12 maanden de plasmaniveaus van de gesuppleerde nutriënten, voorkwam gewichtsverlies en verminderingen in fysieke activiteit en algemene gezondheid, en was kosteneffectief. Een verbeterde fysieke capaciteit na de longrevalidatie bleef behouden voor zowel de interventiegroep als de controlegroep.

**Hoofdstuk 3** onderzoekt of verschillende vormen van motivatie voor het naleven van een gezonde leefstijl daadwerkelijk gerelateerd zijn aan een gezonde leefstijl (fysieke activiteit en een gezond dieet) tijdens en na afloop van een poliklinisch longrevalidatieprogramma. Dit werd onderzocht in het kader van de NUTRAIN trial. Niveaus van geïdentificeerde motivatie (m.a.w. een actie uitvoeren omdat die aansluit bij iemands persoonlijke waarden) waren gerelateerd aan de verandering in de kwaliteit van het dagelijkse voedingspatroon tijdens en na afloop van longrevalidatie; niveaus van geïntrojecteerde motivatie (het uitvoeren van een actie uit angst voor de schaamte die gerelateerd is aan het niet uitvoeren ervan)



waren gerelateerd aan de verandering in fysieke activiteit tijdens en na afloop van de revalidatie.

**Hoofdstuk 4** betreft een narratieve review over de bestaande literatuur over verminderd cognitief functioneren in patiënten met COPD. Dit hoofdstuk toont aan dat enkele ziekte-specifieke factoren, comorbiditeiten en een ongezonde leefstijl kunnen bijdragen tot het ontstaan van cognitieve beperkingen in COPD. Deze beperkingen treffen ongeveer een op de drie patiënten. Veel verschillende cognitieve domeinen kunnen aangedaan worden maar er bestaat geen ziekte-specifiek patroon zoals het geheugenverlies dat karakteristiek is voor de vroege stadia van de ziekte van Alzheimer. Cognitieve beperkingen hebben een negatief effect op de gezondheid, het vermogen tot zelfmanagement, het dagelijks leven en de kwaliteit van leven van patiënten met COPD. Bovendien verhogen cognitieve beperkingen het risico op een ziekenhuisopname.

Voortbouwend op deze review is in **Hoofdstuk 5** onderzocht of de aanwezigheid van het metabool syndroom of sarcopenie gerelateerd is aan een hogere prevalentie en/of een verschillend patroon van cognitieve beperkingen in patiënten met COPD die verwezen waren voor longrevalidatie. Deze metabole aandoeningen komen vaker voor in COPD dan in gezonde controles, en we weten uit eerder onderzoek dat beide metabole condities op zichzelf geassocieerd zijn met een verminderd cognitief functioneren. Het metabool syndroom en sarcopenie hadden een hoge prevalentie in onze bestudeerde COPD-populatie (respectievelijk 54.7% en 30.0%), maar ze bleken niet gerelateerd te zijn aan een hogere prevalentie van cognitieve beperkingen. Dit zijn daarom geen bruikbare indicaties voor aanvullende cognitieve screening.

**Hoofdstuk 6** toont de resultaten van de gerandomiseerde, placebo-gecontroleerde Cogtrain-trial. Dit onderzoek bestond uit een 12-weekse werkgeheugentraining, gevolgd door een 12-weeks onderhoudsprogramma, in patiënten met COPD. Het doel van het onderzoek was om verbeteringen te bewerkstelligen en te consolideren in het cognitief functioneren, de leefstijl en de vatbaarheid voor cognitieve stress. Hoewel het functioneren op de getrainde (werkgeheugen-)taken duidelijk verbeterde, waren er geen verbeteringen te zien op andere cognitieve domeinen na de training. De interventie had ook geen effecten op de andere onderzochte parameters. Werkgeheugentraining lijkt dus geen effectieve manier om het cognitief functioneren en daarmee samenhangende leefstijlaspecten van patiënten met COPD te verbeteren.

**Hoofdstuk 7** plaatst de resultaten van het onderzoek in een breder wetenschappelijk en klinisch perspectief, met een focus op andere benaderingen

## Samenvatting

om het cognitief functioneren en een bestendige gezondere leefstijl te verbeteren bij patiënten met COPD.





**Impact**

Chronic obstructive pulmonary disease (COPD) is one of the highest-burden chronic diseases worldwide. Lung function impairment is characterized by airflow limitation that is usually progressive and largely irreversible (1). COPD is a substantial cause of disability and mortality: 299 million people worldwide were estimated to suffer from the disease in 2017 and COPD was responsible for 3 million deaths in 2016, making it the third most common cause of death worldwide (2).

Although COPD is primarily a lung disease, various manifestations outside the respiratory system are common, including metabolic abnormalities (3, 4), muscle dysfunction (5) and cognitive impairment (CI) (6). CI worsens patients' quality of life (7, 8) and health outcomes (9, 10), it decreases their ability to engage in activities of daily living (11) and it increases their risk of hospitalization (12). All of this contributes to a vicious cycle in which patients' health status deteriorates even further, and to a considerable economic burden for society (13, 14). It is therefore very important to optimally manage COPD from a holistic perspective, consider CI as comorbidity and to assist patients in attaining and maintaining a healthy lifestyle. This thesis investigated determinants of cognitive performance in patients with COPD and the potential efficacy of working memory training (WMT) and nutritional management next to motivational regulations in increasing patients' adherence to a healthier lifestyle.

## Relevance for patients

The results of this thesis are very relevant to patients with COPD. Awareness regarding the adverse impact of their disease and leads for improvement might translate into their health status and wellbeing. Improved prospective memory could for example improve self-management and treatment adherence. Improved executive functioning could enhance self-control, enabling patients to resist tempting unhealthy lifestyle choices in favor of more controlled and healthier ones. The WMT that we developed and evaluated as single intervention was unable to improve overall cognitive functioning or adherence to a healthy lifestyle. Nevertheless, interventional modalities, such as daily-life cognitive stimulation (e.g., playing chess (15), completing sudokus (16) or engaging in social interaction (17, 18)) physical activity (19, 20) and dietary improvement (21, 22), have been shown to benefit cognitive performance in non-COPD populations and are recommended, also in light of other common comorbidities in COPD.

## Relevance for clinical practice

Raising awareness for cognitive impairment in COPD and interventions (also) targeting cognitive functioning in these patients is very relevant for caregivers. The prevalence of CI was particularly high in patients referred for pulmonary rehabilitation (PR) compared to earlier prevalence estimates in stable patients in population studies (6). This, and the fact that CI has been related to adverse health outcomes including the likelihood of dropping out of PR (23), underlines the importance for PR providers to recognize the possible presence of CI in their patients and to tailor educational programs to patients' cognitive capacities.

Furthermore, new insights in specific motivational regulations associated with indicators of adherence to a healthy lifestyle (i.e., physical activity and dietary quality) are important to consider in refining COPD-specific lifestyle counseling interventions.

## Relevance for society

A main aim of this thesis was to determine the health and lifestyle benefits of the investigated interventions. WMT was ineffective, but targeted nutritional management improved patients' health status at an increase of costs that might be considered acceptable for patients with a high disease burden. Moreover, motivational interventions, if proven feasible as discussed above, might cost-effectively improve patients' lifestyle, health status, quality of life and functional capacities as well. This enables patients to live more meaningful lives for a longer period of time, for instance by prolonging their ability to work. Furthermore, it might decrease health care expenditure and the COPD-associated disability burden. Both are relevant for society, as rising health care costs present a large and growing economic burden and COPD is the fifth-leading cause of disability-adjusted life years (DALYs) worldwide (24).

## Relevance for commercial parties

Motivational counselling does not have to remain confined to health care professionals. In fact, counselling by commercial parties would take work away from health care professionals, which would alleviate the growing work pressure and financial burden on the health care system.

Incorporating counselling into an e-health application could reduce the workload associated with counselling even further. Various applications have been developed over the past years to improve chronic disease management with promising results on health outcomes, such as myIBDcoach (25, 26) for inflammatory bowel disease and SanaCoach COPD for COPD (27). Counselling could even take place entirely virtually. Haptic, an app developed to improve the lifestyle of corporate employees, already uses such virtual, yet highly realistic, avatars.

The NUTRAIN trial showed that nutritional supplementation can improve the overall health status of patients with COPD in a cost-effective manner. Further developing medical foods and/or supplements, aimed at improving physical capacity and health status but also building forth upon promising results regarding their benefits on cognitive functioning (28, 29), for instance, could therefore be promising for commercial parties.

Developing WMT modalities and marketing them as games has been an important focus of industry in the past 20 years. Although these games are often supported by claims that their effectiveness has been proven, independent investigations routinely indicate they do not improve cognitive or real-life task performance (30-32). The results of the current thesis also indicate that further developing WMT applications is likely not worthwhile.



## References

1. Global Initiative for Chronic Obstructive Lung Disease (GOLD). Global Strategy for the Diagnosis, Management and Prevention of Chronic Obstructive Pulmonary Disease. Agustí A, Vogelmeier C, editors; 2020.
2. World Health Organization. The top 10 causes of death 2018 [accessed July 16, 2020]. Available from: <https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death>.
3. Trajanoska K, Schoufour JD, Darweesh SKL, Benz E, Medina-Gomez C, Alferink LJM, *et al.* Sarcopenia and its clinical correlates in the general population: the Rotterdam Study. *Journal of Bone and Mineral Research*. 2018;33(7):1209-18.
4. Cebron Lipovec N, Beijers RJ, Van den Borst B, Doehner W, Lainscak M, Schols AMWJ. The prevalence of metabolic syndrome in chronic obstructive pulmonary disease: a systematic review. *COPD*. 2016;13(3):399-406.
5. Remels AH, Gosker HR, Langen RC, Schols AMWJ. The mechanisms of cachexia underlying muscle dysfunction in COPD. *Journal of Applied Physiology*. 2013;114(9):1253-62.
6. Yohannes AM, Chen W, Moga AM, Leroi I, Connolly MJ. Cognitive impairment in chronic obstructive pulmonary disease and chronic heart failure: a systematic review and meta-analysis of observational studies. *JAMDA*. 2017;18:451.e1-.e11.
7. Dodd JW, Charlton RA, Van den Broek MD, Jones PW. Cognitive dysfunction in patients hospitalized with acute exacerbation of COPD. *Chest*. 2013;144(1):119-27.
8. Roncero C, Campuzano AI, Quintano JA, Molina J, Perez J, Miravittles M. Cognitive status among patients with chronic obstructive pulmonary disease. *International Journal of COPD*. 2016;11:543-51.
9. Brega AG, Grigsby J, Kooken R, Hamman RF, Baxter J. The impact of executive cognitive functioning on rates of smoking cessation in the San Luis Valley Health and Aging Study. *Age and Ageing*. 2008;37(5):521-5.
10. Meek P, Lareau SC, Anderson D. Memory for symptoms in COPD patients: how accurate are their reports? *European Respiratory Journal*. 2001;18(3):474-81.
11. Baird C, Lovell J, Johnson M, Shiell K, Ibrahim JE. The impact of cognitive impairment on self-management in chronic obstructive pulmonary disease: a systematic review. *Respiratory Medicine*. 2017;129:130-9.
12. Chang SS, Chen S, McAvay GJ, Tinetti ME. Effect of coexisting chronic obstructive pulmonary disease and cognitive impairment on health outcomes in older adults. *Journal of the American Geriatric Society*. 2012;60(10):1839-46.
13. Mannino DM, Thorn D, Swensen A, Holguin F. Prevalence and outcomes of diabetes, hypertension and cardiovascular disease in COPD. *European Respiratory Journal*. 2008;32(4):962-9.

14. Mannino DM, Higuchi K, Yu T-C, Zhou H, Li Y, Tian H, et al. Economic burden of COPD in the presence of comorbidities. *Chest*. 2015;148(1):138-50.
15. Lillo-Crespo M, Forner-Ruiz M, Riquelme-Galindo J, Ruiz-Fernández D, García-Sanjuan S. Chess practice as a protective factor in dementia. *International Journal of Environmental Research and Public Health*. 2019;16(12):2116.
16. Ferreira N, Owen A, Mohan A, Corbett A, Ballard C. Associations between cognitively stimulating leisure activities, cognitive function and age-related cognitive decline. *International Journal of Geriatric Psychiatry*. 2015;30(4):422-30.
17. Kelly ME, Duff H, Kelly S, McHugh Power JE, Brennan S, Lawlor BA, et al. The impact of social activities, social networks, social support and social relationships on the cognitive functioning of healthy older adults: a systematic review. *Systematic Reviews*. 2017;6:259.
18. Biddle KD, D'Oleire Uquillas F, Jacobs HIL, Zide B, Kirn DR, Rentz DM, et al. Social engagement and amyloid- $\beta$ -related cognitive decline in cognitively normal older adults. *American Journal of Geriatric Psychiatry*. 2019;27(11):1247-56.
19. Falck RS, Davis JC, Best JR, Crockett RA, Liu-Ambrose T. Impact of exercise training on physical and cognitive function among older adults: a systematic review and meta-analysis. *Neurobiology of Aging*. 2019;79:119-30.
20. Song D, Yu DSF, Li PWC, Lei Y. The effectiveness of physical exercise on cognitive and psychological outcomes in individuals with mild cognitive impairment: a systematic review and meta-analysis. *International Journal of Nursing Studies*. 2018;79:155-64.
21. Martínez-Lapiscina EH, Clavero P, Toledo E, Estruch R, Salas-Salvadó J, San Julián B, et al. Mediterranean diet improves cognition: the PREDIMED-NAVARRA randomised trial. *Journal of Neurology, Neurosurgery and Psychiatry*. 2013;84(12):1318-25.
22. Van den Brink AC, Brouwer-Brolsma EM, Berendsen AAM, Van de Rest O. The Mediterranean, Dietary Approaches to Stop Hypertension (DASH), and Mediterranean-DASH Intervention for Neurodegenerative Delay (MIND) diets are associated with less cognitive decline and a lower risk of Alzheimer's disease - a review. *Advances in Nutrition*. 2019;10(6):1040-65.
23. Cleutjens FAHM, Spruit MA, Ponds RWHM, Vanfleteren LEGW, Franssen FM, Dijkstra JB, et al. The impact of cognitive impairment on efficacy of pulmonary rehabilitation in patients with COPD. *JAMDA*. 2017;18(5):420-6.
24. GBD 2017 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392(10159):1789-858.

25. De Jong M, Van der Meulen-De Jong A, Romberg-Camps M, Becx M, Maljaars JP, Cilissen M, *et al.* Telemedicine for management of inflammatory bowel disease (myIBDcoach): a pragmatic, multicentre, randomised controlled trial. *Lancet*. 2017;390(10098):959-68.
26. De Jong M, Van der Meulen-De Jong A, Romberg-Camps M, Degens J, Becx M, Markus T, *et al.* Development and feasibility study of a telemedicine tool for all patients with IBD: MyIBDcoach. *Inflammatory Bowel Disease*. 2017;23(4):485-93.
27. Trappenburg J, Niesink A, De Weert-Van Oene GH, Van der Zeijden H, Van Snippenburg R, Peters A, *et al.* Effects of telemonitoring in patients with chronic obstructive pulmonary disease. *Telemedicine Journal and E-Health*. 2008;14(2):138-46.
28. Külzow N, Witte AV, Kerti L, Grittner U, Schuchardt JP, Hahn A, *et al.* Impact of omega-3 fatty acid supplementation on memory functions in healthy older adults. *Journal of Alzheimer's disease*. 2016;51(3):713-25.
29. Troesch B, Eggersdorfer M, Laviano A, Rolland Y, Smith AD, Warnke I, *et al.* Expert opinion on benefits of long-chain omega-3 fatty acids (DHA and EPA) in aging and clinical nutrition. *Nutrients*. 2020;12(9):2555.
30. Etherton JL, Oberle CD, Rhoton J, Ney A. Effects of Cogmed working memory training on cognitive performance. *Psychological Research*. 2019;83(7):1506-18.
31. Harris DJ, Wilson MR, Vine SJ. A systematic review of commercial cognitive training devices: implications for use in sport. *Frontiers in Psychology*. 2018;9:709.
32. Kable JW, Caulfield MK, Falcone M, McConnell M, Bernardo L, Parthasarathi T, *et al.* No effect of commercial cognitive training on brain activity, choice behavior, or cognitive performance. *Journal of Neuroscience*. 2017;37(31):7390-402.



# Dankwoord

Ik zie mezelf nog bij Juul in de tuin staan op het moment dat ik een promotieplek in Leuven aangeboden kreeg. Die positie zei ik af, ik ging weer een paar maanden in de supermarkt werken, en daarop kreeg ik te horen "dat ik drie jaar zonder baan zou zitten" en ik "er nooit meer tussen zou komen". Inmiddels kunnen we denk ik stellen dat die conclusies enigszins voorbarig waren.

Mijn masterscriptiebegeleidster vatte in haar dankwoord haar promotietraject samen in één uitspraak. "Het was vaak fantastisch en soms verschrikkelijk". Dat wil ik graag volmondig beamen. Het was fantastisch wanneer gezellige proefpersonen langskwamen voor meetdagen, wanneer ik borrels en symposia organiseerde met mijn 'vriendenteam' van de aio-raad, of wanneer ik me staande hield presenterend in een zaal vol professoren op een groot congres. Het was verschrikkelijk op de nachten waarop ik om vijf uur wakker werd, alles voor me zag wat fout kon gaan met mijn onderzoek, en vervolgens vier wakkere uren later met kleine oogjes achter mijn bureau schoof. Hierbij wil ik graag de belangrijkste mensen bedanken die er de afgelopen vier jaar bij waren.

Annemie, ik ben ontzettend dankbaar voor de kans die je me gaf om te promoveren. Het was echt niet vanzelfsprekend om deze promotieplek te geven aan een 22-jarig jochie zonder enige achtergrond in longziekten. Ik heb altijd ontzettend veel bewondering gehad voor het gruwelijk hoge niveau waarop je denkt en heb enorm genoten van de mogelijkheid om daarvan te leren. Ik moet je bij tijden gek gemaakt hebben met al mijn vlaaipunten, maar ik hoop dat deze promotie je vertrouwen in me terugbetaalt.

Harry, jouw bijdrage aan mijn promotietraject wordt perfect belichaamd door je reddingsactie op de Maas, toen jij en Ramon de *banter boat* met mij en Justine naar het eindpunt gidsten. Drive en enthousiasme heb ik altijd genoeg gehad, maar aan richting ontbrak het wel eens, en die richting heb je zowel op de Maas als in mijn promotietraject verzorgd. Bedankt voor alle keren dat je me wees op dingen die ik niet handig aanpakte, dat je me terechtwees als ik om de zaken heen draaide als ze niet gingen zoals ze moesten gaan - en uiteraard voor die reddingsactie op de Maas.

Zonder al mijn proefpersonen was dit proefschrift er natuurlijk ook niet geweest. U heeft de moeite genomen om een aantal tientallen uren van uw leven naar een computerschermpje te turen, bijna de helft van u zelfs in de placebogroep. Hoewel het niet altijd even leuk was (dat heb ik van sommigen van u wel begrepen), hoop ik dat de gezelligheid tijdens de afspraken op de universiteit daar wat voor gecompenseerd heeft. Het patiëntencontact is voor mij in ieder geval erg waardevol geweest. Voor de gezelligheid, uiteraard, maar nog veel meer om een

beetje een idee te krijgen van wat er achter de woorden en statistieken in dit boekje zit.

Mijn kamergenoten waren in meerdere opzichten mijn reddingsboei. Stephanie in het prille begin bij MRI-meetdagen en METC-aanvragen, en om me met zachte hand wat realisme in te praten als mijn planning weer eens wat te optimistisch bleek. Ik heb het je in de afgelopen jaren al vaker gezegd, maar wat moet ik je bij tijden tot wanhoop hebben gedreven. Lianne als grote ondersteuning binnen Cogtrain en om me er regelmatig aan te herinneren dat ik soms eens wat minder slordig moest zijn. En Vasili: kamergenoot, vriend, paranimf, slapie, drinkmaatje, wingman, relatie-adviseur, en degene die binnen no-time een straalkacheltje geregeld kreeg als de verwarming er thuis weer eens mee ophield. Reddingsboei doet de beschrijving van je rol tekort.

Er komt natuurlijk maar één andere persoon in aanmerking om mijn paranimf te zijn. Viv, wij zijn er al jaren altijd voor elkaar, niet in het minst tijdens onze wandelingetjes door Heugem en De Heeg, of tijdens onze dinertjes. Bij mij thuis, in Huize Mimosa of sinds kort in Helden-Panningen. Ik ben enorm trots op je, super blij dat je zo goed op je plek zit, en ik hoop nog vele foto's met foute Snapchat-filters met je te kunnen maken.

De collega's met wie ik ontelbare koffietjes, lunches, dinertjes in het personeelsrestaurant en avonden clubpower gedeeld heb, in het bijzonder Inez, Ellen, Denise, Philippe, Timme, Justine en Sara.

Mijn collega's van de aio-raad: Annet, Charlotte, Elisa, Eva, George, Gianluca, Heike, Jacco, Kelly, Mattea, Mirjam, Pauline en Sara.

Mijn andere huidige en voormalige collega's van pulmonologie: Anita, Ardy, Bregtje, Caspar, Cheryl, Chiel, Coby, Judith, Jules, Juliette, Lieke, Karin, Marco, Mieke, Niki, Pieter, Ramon, Rianne, Rosanne, Sara(h), Sophie, Viviënne, Wessel, Wouter G en Wouter v/d W.

Hurra, hurra, die V2E is auch noch da! Hangry buien in Gent, het monster van Toth in Antwerpen, de grote triomfen en diepe dalen van ons voetbalcluppie die we door het hele land heen hebben gezien: jullie zijn al bijna vijftien jaar lang een constante factor. Het feit dat ik de eerste maar zeker niet de laatste promovendus van ons groepje ben is hopelijk het ultieme bewijs van onze VWO+-mentaliteit...

Pa, voor alle PSV-gerelateerde hoogte- en dieptepunten die we de afgelopen twintig jaar hebben gedeeld. Tien jaar lang op vak UU, rij 37, stoel 47; tegenwoordig thuis op de bank. Het is fijn dat jij mijn innerlijke voetbaltrainer

een beetje kan kanaliseren, en dat we 't dan ten minste eens zijn over dat iemand als Pablo Rosario niet bij een club als PSV in de basis hoort.

En als laatste natuurlijk de belangrijkste vrouwen in mijn leven. Fré, omdat je er altijd, altijd voor me bent. Ik kan niet wachten om met je samen te gaan wonen en samen een mooie toekomst (verder) op te bouwen. Juul, voor de driekwartliterflessen La Chouffe die we leeg hebben getutterd, en mamma - misschien dat ik wel het meeste zeg als ik zeg dat jouw bijdrage aan mijn promotie en mijn ontwikkeling in het algemeen niet in een dankwoord te beschrijven is. Wie weet waar ik was geweest zonder jouw hulp en advies, en zonder onze wandelrondjes langs het kanaal. In ieder geval niet hier in de positie waarin ik je kan bedanken voor alles wat je voor me doet en gedaan hebt.







# Curriculum vitae

Martijn van Beers was born on June 3rd, 1994, in Veldhoven, The Netherlands. He followed preparatory scientific education (VWO) at the 'Heerbeeck College' secondary school in Best (Noord-Brabant, The Netherlands). After graduating in 2012, he successfully completed the bachelor program in Psychology at Tilburg University in 2015, where he wrote his thesis under the supervision of dr. Ruth Mark. Subsequently, he obtained a master in Neuropsychology at Maastricht University in 2016. He wrote his master thesis at Hasselt University (Belgium) under the supervision of dr. Ellen Jongen en dr. Judith Urlings.

In December 2016 he started his PhD research at the Department of Respiratory Medicine at Maastricht University, where he conducted the research described in this thesis, investigating determinants of cognitive performance in patients with chronic obstructive pulmonary disease and the potential efficacy of working memory training next to motivational regulations in increasing patients' adherence to a healthy lifestyle. He performed this research under the supervision of prof. dr. ir. Annemie Schols and dr. Harry Gosker. During this period he also presented his work at various national and international conferences.





# List of publications

**Van Beers M**, Janssen DJA, Gosker HR, Schols AMWJ. Cognitive impairment in chronic obstructive pulmonary disease: disease burden, determinants and possible future interventions. *Expert Review of Respiratory Medicine*. 2018;12(12):1061-74.

**Van Beers M**, Rutten-Van Mölken MPMH, Van de Bool C, Boland M, Kremers SPJ, Franssen FME, Van Helvoort A, Gosker HR, Wouters EF, Schols AMWJ. Clinical outcome and cost-effectiveness of a 1-year nutritional intervention programme in COPD patients with low muscle mass: the randomized controlled NUTRAIN trial. *Clinical Nutrition*. 2020;39(2):405-13.

**Van Beers M**, Gosker HR, Janssen DJA, Cleutjens FAHM, Franssen FME, Van Bortel MPJ, Wouters EF, Ponds RWHM, Schols AMWJ. Cognitive performance in relation to metabolic disturbances in patients with COPD. *Clinical Nutrition*; in press.

**Van Beers M**, Meis JJM, Van de Bool C, Schols AMWJ, Kremers SPJ. Motivational regulations and lifestyle in patients with COPD and low muscle mass before and after pulmonary rehabilitation. Manuscript submitted.

**Van Beers M**, Mount SW, Houben K, Gosker HR, Schuurman L, Franssen FME, Janssen DJA, Schols AMWJ. Efficacy of working memory training in chronic obstructive pulmonary disease: the randomized placebo-controlled Cogtrain trial. Manuscript submitted.